


Enhancing Coordination of Behavioral Health Services after Superstorm Sandy: Planning for Future Disasters



Final Data Profile: Atlantic County Medicare Fee-for-Service Beneficiaries

*Demographics, Behavioral Health
Conditions, and Utilization of
Health Services*

June 17, 2014

This page left intentionally blank

TABLE OF CONTENTS

Preface	1
Introduction	2
HQSI Project Team	7
Acknowledgements	8
Executive Summary	9
Demographics	11
• Total Medicare FFS Beneficiary Population by County	11
• Percent of Medicare FFS Beneficiaries in the General Population	12
• Percent of Medicare FFS Beneficiary Population by Gender by County	12
• Percent of Medicare FFS Beneficiary Population by Race by County.....	13
• Percent of Medicare FFS Beneficiary Population by Age by County	14
• Income Status by County	15
Behavioral Health Conditions	17
• Prevalence and Incidence.....	17
– Summary	18
– Depression or Proxy Disorders	21
– Depression.....	27
– Anxiety Disorders	28
– Adjustment Disorders	29
– Post-Traumatic Stress Disorder.....	30
– Alcohol or Substance Abuse	31
– Substance Abuse Alone	32
– Suicide and Intentional Self-Inflicted Injury	33
• Risk Factors for Depression or Proxy Disorders.....	34
– Summary	34
– Any of the Top Five Risk Factors for Depression or Proxy Disorders	35
– Alzheimer’s Disease and Related Disorders or Senile Dementia.....	38
– Sleep Disturbance.....	38
– Substance or Alcohol Abuse or Tobacco Use	39
– Hip/Pelvic Fractures	39
– Amputations	40
Utilization	41
• Outpatient Behavioral Health Services	41
– Assessments	41
› Summary	41
› Depression Screening	43
› Diagnostic Psychological Tests	47

TABLE OF CONTENTS

› Health and Behavior Assessment/Intervention.....	48
› Neuropsychological Tests.....	49
› Psychiatric Diagnostic Procedures.....	51
– Therapies	53
› Summary	53
› Individual Psychotherapy.....	55
› Family Psychotherapy.....	57
› Group Psychotherapy.....	58
› Electroconvulsive Therapy	59
› Biofeedback Therapy.....	60
• Inpatient Services.....	61
– Summary.....	61
– Psychiatric Hospital Admissions.....	63
– Acute Care Hospitals	64
› Admissions.....	64
› Observation Stays.....	65
› Emergency Department Visits.....	66
– Within 30 Days of Acute Care Hospital Discharge	67
› Summary	67
› 30-Day Hospital Readmissions.....	69
› Observation Stays Within 30 Days of Discharge	70
› Emergency Department Visits Within 30 Days of Discharge	71
– Other Settings.....	72
› Summary	72
› Home Health Agency Services	74
› Skilled Nursing Facility Services	75
› Hospice Services.....	76
› Medical Rehabilitation Services	77
Appendices.....	79
• Appendix A: Behavioral Health Conditions.....	79
• Appendix B: Risk Factors for Depression or Proxy Disorders.....	81
• Appendix C: Utilization of Outpatient Mental Health Services	85
• Appendix D: Utilization of Services – Inpatient and Other Settings	87
• Appendix E: Time Frames and Formulae	89
• Appendix F: Professional Type by Behavioral Health Services.....	90
• Appendix G: References.....	91
Index of Figures	93

PREFACE

On October 29, 2012, Superstorm Sandy hit the Eastern Seaboard, impacting more than a dozen states. New Jersey, which took the brunt of the storm along its densely populated coastline, was devastated. Thousands of residents were displaced, their homes and communities damaged or destroyed.

Lessons learned from prior natural disasters showed that victims of storms like Superstorm Sandy are often at an elevated risk for behavioral health issues such as post-traumatic stress disorder (PTSD), depression, and substance abuse.^{1,2} While disaster-related issues subside over time, evidence shows that victims can experience a prolonged period of elevated risk, especially those with pre-existing mental health issues.³ Older adults and disabled residents with mental health conditions are at increased risk of deteriorating health, depression, increased isolation, and breakdown in the continuum of health care. Additionally, past natural disasters also show that access to informational resources on disaster-related mental health disorders, outcomes, and service utilization are important factors to consider.^{4,5}

This final county profile can help healthcare professionals learn more about the behavioral health status and healthcare utilization patterns of Medicare Fee-for-Service (FFS) beneficiaries before and after Superstorm Sandy. As such, it may be a useful tool in planning for future disasters. This profile is one of 10 created for each of the Federal Emergency Management Agency (FEMA)-declared disaster counties in New Jersey. The profiles explore county-level health status and health determinants of post-disaster spikes in behavioral health issues and treatments. This last update includes one more quarter of comprehensive post-Sandy data than the previous profile, which was published in May 2014.

INTRODUCTION

Enhancing Coordination of Behavioral Health Services after Superstorm Sandy: Planning for Future Disasters is a Special Innovation Project funded by the Centers for Medicare & Medicaid Services (CMS). As part of this project, Healthcare Quality Strategies, Inc. (HQSI), the quality improvement organization (QIO) for New Jersey, studied data on prevalence and incidence of selected behavioral health conditions, the utilization of health services, and demographic information from the Medicare claims for Medicare FFS beneficiaries residing in the 10 New Jersey FEMA-declared disaster counties after Superstorm Sandy. These counties include Atlantic, Bergen, Cape May, Essex, Hudson, Ocean, Middlesex, Monmouth, Somerset, and Union.

From its analysis, HQSI created data profiles for each of these FEMA-designated counties. The initial set of county profiles, which covered the period January 1, 2011 to March 31, 2013, was published in January 2014. These profiles were then updated in May 2014 and covered the period from January 1, 2011 to September 30, 2013. This final profile is the last update planned for Atlantic County and includes data from January 1, 2011 to December 31, 2013. This profile can be used to determine and compare the prevalence and incidence of the selected behavioral health conditions and utilization of services among all 10 FEMA-declared disaster counties before and after Superstorm Sandy.

HQSI also created profiles for a subset of 10 communities. These communities were selected because they had high rates of Medicare FFS beneficiaries both with and at risk for depression or proxy disorders and other factors. The initial community profiles, along with the first updated version, are available at www.hqsi.org. The community profiles can be used to determine and compare the prevalence and incidence of the selected behavioral health conditions and utilization of services in the selected communities compared to their counties.

The county and community profiles are based on Medicare FFS claims data and provide a glimpse into the prevalence and incidence of selected behavioral health conditions and risk factors for depression, as well as the utilization of Medicare-covered behavioral health services among Medicare beneficiaries residing in the selected counties or communities before and after Superstorm Sandy. Since patients with behavioral health conditions may receive other health services because of medical problems caused by their behavioral health conditions, or may avoid utilizing behavioral health services, this profile also looks at the utilization of non-behavioral health services.

These profiles are being shared with state and local governments and agencies, health care providers, community-based organizations, and the research community to support a community-based approach to enhance the coordination of behavioral health services after a natural disaster, and to increase utilization of the Medicare depression screening benefit which became a covered service in October 2011.

WHAT'S NEW IN THIS UPDATE

This second updated profile shows four quarters of post-Sandy data, with the most updated claims from January 2011 to December 2013. This profile compares the 12-month rates from the year before and after the storm. In this profile, we reference October 2011 to September 2012 as the year before Superstorm Sandy and January 2013 to December 2013 as the year after the storm.

HOW TO USE THIS PROFILE

This profile includes an analysis of the eight behavioral health conditions which, based on literature review and feedback from the subject matter experts consulted for this project, were found to increase after natural disasters.

This profile is divided into the following sections, each of which is preceded by a user-friendly overview:

- Demographics (page 11)
- Prevalence and incidence of behavioral health conditions (page 17)
- Risk factors for depression or proxy disorders (page 34)
- Utilization of outpatient behavioral health assessments (page 41)
- Utilization of outpatient behavioral health therapies (page 53)
- Utilization of inpatient health services (page 61)
- Utilization of inpatient health services within 30 days of discharge (page 67)
- Utilization of other settings (page 72)

Here are some additional tips for using this profile:

- Use the Executive Summary (pages 9-10) for a quick overview of this profile's key points, as well as a snapshot table that summarizes the prevalence of the selected behavioral health conditions and utilization of behavioral health services before and after Sandy
- Use the Behavioral Health Conditions section (pages 17-33) for in-depth analyses and graphical comparison on the prevalence and incidence of eight behavioral health conditions before and after Superstorm Sandy
- Use the New Jersey and county maps to: identify areas with higher rates of Medicare FFS beneficiaries at risk for depression and proxy disorders (pages 25-26); and areas with low utilization of the depression screening benefit (pages 45-46)

METHODOLOGY

Each county profile compares one county's statistics to the aggregate of the 10 counties and to the other nine counties. Primary data sources include Medicare FFS Part A and Part B claims, the Medicare enrollment database, and U.S. Census data. The Medicare enrollment database includes basic demographic statistics such as age, gender, and race while the U.S. Census data provides a proxy indicator (average household income) for socio-economic status.⁶ Based on the ICD-9-CM (International Classification of Disease, Ninth Revision, Clinical Modification), CPT (Current Procedural Terminology) or HCPCS (Healthcare Common Procedure Coding System) codes in Medicare Part A and Part B claims, beneficiaries were identified for diseases/conditions related to behavioral health conditions such as depression. Appendices A through G contain documentation, technical notes, codes, algorithms, data sources, and references.

Medicare Part A claims were also used to analyze utilization of health services in acute care hospitals, skilled nursing facilities, medical rehabilitation facilities, home health agencies, hospice, and inpatient psychiatric facilities. Medicare Part A and Part B claims provide information on the utilization of mental health outpatient services for assessment (e.g., depression screening, diagnostic psychological tests) and treatment (e.g., individual psychotherapy, biofeedback therapy).

To identify beneficiaries with an elevated risk of depression after the storm, HQSI conducted a literature review of risk factors for depression (see Appendix B). Previous studies identified psychosocial and biological factors, increased age, history of cancer, Parkinson's disease, Alzheimer's disease, changes in mental function, and medication side effects as risk factors for developing depression. Based on findings from the literature review and factors available through Medicare claims, logistic regression analysis was conducted with Medicare claims, and the top five risk factors (Alzheimer's disease and related disorders or senile dementia, hip/pelvic fractures, amputations, substance or alcohol abuse or tobacco use, and sleep disturbance) were used to identify beneficiaries with high risk for developing depression or proxy disorders (i.e., anxiety and adjustment disorders).

MEASUREMENT TIME FRAMES

This profile includes data from January 1, 2011 through December 31, 2013. Results are presented using different charts and measurement time frames as follows:

- Annual bar charts show the annual rates in the year before (October 1, 2011 to September 30, 2012) and after (January 1, 2011 to December 31, 2013) Superstorm Sandy. Statistics on demographics, prevalence of behavioral health conditions, and utilization of health services are presented for this 12-month period. These statistics allow for comparison across affected counties before and after Superstorm Sandy
- Annual trend charts with rolling quarters for the behavioral health conditions and utilization statistics are included to adjust for seasonal variation and to examine possible changes in the year before and after Superstorm Sandy. The time period includes nine data points from January 1, 2011 to December 31, 2013
- Annual percent change (relative change) bar charts show relative increase or decrease in rates from the year before and after Superstorm Sandy. These statistics allow for comparison across the 10 affected counties and to analyze the potential impact of Superstorm Sandy
- Quarterly new incidence charts for eight behavioral health conditions include eight quarters of data from January 1, 2012 to December 31, 2013. This allows for the identification of new cases in a given quarter when compared to the prior year
- Quarterly line charts show the trend in the utilization of depression screening for eight quarters from January 1, 2012 to December 31, 2013.

DATA CONSIDERATIONS

There are now four quarters of post-storm data available, which is reflected in this final update. The claims data processing lag of at least six months, coupled with the one-year project time frame, reduces the optimal time frame for more accurate estimation of post-Sandy effects.

Identification of beneficiaries with behavioral health conditions is based on diagnoses being reported in Medicare FFS claims and could result in underestimation. There is no accurate way to identify when certain health conditions began and ended when claims data is used.

According to the subject matter experts consulted for this project, unlike other conditions, behavioral health issues are often underdiagnosed in our society and the stigma associated with behavioral health conditions may prevent people from seeking care in mental health facilities. The subject matter experts also indicated that estimating the prevalence of depression using claims data can be particularly difficult as depression is often undiagnosed or not documented. Depression can be present with symptoms of anxiety and adjustment disorders. Based on this feedback, a combination measure named “depression or proxy disorders” was created to estimate prevalence and incidence of depression. If a patient has at least one of the three conditions reported in Medicare claims, he/she will be flagged as having depression or proxy disorders.

This county profile can be used to compare the prevalence and incidence rates of eight selected behavioral health conditions (see page 19) based on the ICD-9-CM codes through the analysis of Medicare claims. This profile may be used to prioritize and plan community and county preparation for the care, tracking, and monitoring of Medicare beneficiary behavioral health status and health care utilization patterns.

This is the final update of these data profiles which includes one more quarter of data than the previous profile during the post-Superstorm Sandy time period as the project ends on July 31, 2014.

Martin P. Margolies

Chief Executive Officer

Mary Jane Brubaker, MCIS

Chief Operating Officer

Diane Babuin, MS, CPHQ

Director, Quality Improvement and Communications

Ya-ping Su, PhD

Director, Research and Analysis

Suzanne Dalton, RN, BS, EdM

Project Manager

Andrew Miller, MD, MPH

Medical Director

Mona Abdalla, BA

Administrative Associate

Christine Aisenberg, BA

Proofreader

Kathy Brown, BS

InDesign Specialist

Zhengyu Bu, MS

Health Services Research Analyst

Sue Chen, MS

Statistician

Wei-Yi Chung, MS

Database Administrator

Barbara Coleman

Administrative Associate

Dawn Cullen, BA

Communications Specialist

Ashley Dopp, BA

Communications Specialist

Karen Hale, MEd

Community Liaison

Kim Karnell, BS

Information Specialist

Janet Knoth, BS, RN, CHPN, CPHQ

Quality Improvement Specialist

Judy Miller, MS, RN

Quality Improvement Specialist

Olubukunola Oyedele, MPH

Community Liaison

Rita Pascale

Administrative Associate

Barbara Perzyna, BS

Visual Communications Specialist

Ziphora Sam, MPH

Epidemiologist

Marianne Sagarese, BSN, RN

Quality Improvement Specialist

Nicole Skyer-Brandwene, MS, RPh, BCPS

Quality Improvement Specialist

ACKNOWLEDGEMENTS

Special thanks to the subject matter experts who assisted with the project by providing feedback and guidance to the HQSI project team.

Carol Benevy, MSW

New Jersey Hope and Healing Project
Barnabas Health Institute for Prevention

Mary Ditri, MA, CHCC

New Jersey Hospital Association

Adrienne Fessler-Belli, MSW, LCSW

New Jersey Department of Human Services
Disaster & Terrorism Branch

Mark Firth, MA, MSW

New Jersey Department of Human Services
Division of Mental Health and Addiction Services

Mary Goepfert, MPA, APR, CPM

New Jersey Group for Access and Integration Needs in Emergencies and Disasters

Sheldon Green

New Jersey Primary Care Association

Connie Greene, MA, CAS, CSW, CPS

Barnabas Health Institute for Prevention

Bob Kley

Mental Health Association in New Jersey, Inc.

Lynn Kovitch, MEd

New Jersey Department of Human Services
Division of Mental Health and Addiction Services

Karen McCoy, RN, BSN

Home Care Association of New Jersey

Elyse Perweiler, MPP, RN

NJ Institute for Successful Aging

Lynn Stefanowicz, MA, LCSW

Meridian Behavioral Health

Megan Sullivan, LPC, LCADC, DRCC

New Jersey Department of Human Services
Disaster & Terrorism Branch

Pete Summers

The New Jersey Association of County and City Health Officials (NJACCHO)

KEY OBSERVATIONS

The following observations show Atlantic County's percent change and ranking among all 10 counties after Superstorm Sandy among Medicare FFS beneficiaries.


1. There was a 2.52% relative increase in depression or proxy disorders, a 4.85% relative increase in anxiety disorders, a 0.03% relative increase in adjustment disorders, and a 3.57% relative increase in PTSD.
2. The highest rates of depression or proxy disorders in Atlantic County were among Hispanic beneficiaries (232.69 per 1,000 beneficiaries), female beneficiaries (251.68 per 1,000 beneficiaries), and beneficiaries below 65 years old (364.82 per 1,000 beneficiaries).
3. Atlantic County had highest rate of suicide and intentional self-inflicted injuries (6.06 per 1,000 Medicare FFS beneficiaries) despite a 6.34% relative decrease in prevalence.
4. There was a 2.39% relative increase in the prevalence of any of the top five risk factors for depression or proxy disorders.
5. There also was a relative increase in the following risk factors: substance or alcohol abuse or tobacco use (3.50%) and sleep disturbances (7.76%).
6. Atlantic County had the lowest rate of hip/pelvic fractures (6.08 per 1,000 beneficiaries) among all 10 counties.
7. Utilization of depression screening increased from 1.12 per 1,000 Medicare FFS beneficiaries to 11.61, representing the highest relative increase among all 10 counties.
8. Atlantic County had the highest utilization rate of health and behavior assessment/intervention (1.41 per 1,000 beneficiaries) among all 10 counties.
9. Atlantic County experienced a relative decrease in the utilization of individual psychotherapy (8.03%) and psychiatric diagnostic procedures (10.70%).
10. Atlantic County had the highest utilization of acute care hospital visits (330.22 per 1,000 beneficiaries), emergency department visits (263.07 per 1,000 beneficiaries), and observation stays (20.83 per 1,000 beneficiaries) among all 10 counties.
11. Atlantic County had the highest utilization of emergency department visits within 30 days of discharge (81.44 per 1,000 beneficiaries) despite a 15.75% relative decrease in the use of this service.
12. Atlantic County had the lowest utilization of skilled nursing facility services (57.91 per 1,000 beneficiaries) among all 10 counties.

This *Snapshot of Atlantic County* summarizes the prevalence of the behavioral health conditions, as well as risk factors for depression or proxy disorders, analyzed for this profile. This *Snapshot* also lists the most frequently performed behavioral health assessments and therapies in Atlantic County compared to the average among all 10 counties. It illustrates the change in conditions and utilization of services before and after Sandy.

Figure 1. Snapshot of Atlantic County						
Behavioral Health Conditions	Prevalence per 1,000 Medicare FFS Beneficiaries					
	Atlantic County			10 County Rate		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change
Depression or Proxy Disorders	204.91	210.08	2.52	192.99	197.65	2.41
• Depression alone	126.73	125.42	-1.03	124.72	125.36	0.51
• Anxiety Disorders alone	122.86	128.82	4.85	105.70	113.91	7.77
• Adjustment Disorders alone	30.28	30.29	0.03	29.82	29.09	-2.45
Alcohol or Substance Abuse	42.98	39.76	-7.49	30.51	33.73	10.55
• Substance Abuse alone	25.26	22.18	-12.19	16.71	17.54	4.97
PTSD	5.32	5.51	3.57	4.18	4.69	12.20
Suicide and Intentional Self-Inflicted Injury	6.47	6.06	-6.34	4.40	4.39	-0.23
Top Five Risk Factors* for Depression or Proxy Disorders						
• Any of the Top Five Risk Factors	145.06	148.52	2.39	136.36	136.15	-0.15
• Substance or Alcohol Abuse or Tobacco Use	89.50	92.63	3.50	78.33	81.78	4.40
• Alzheimer’s Disease and related disorders or Senile Dementia	33.94	33.45	-1.44	39.11	34.91	-10.74
• Sleep Disturbance	26.94	29.03	7.76	24.24	24.78	2.23
• Hip/Pelvic Fractures	6.47	6.08	-6.03	7.95	7.66	-3.65
• Amputations	1.17	1.17	0.00	1.11	0.99	-10.81
Behavioral Health Services	Utilization per 1,000 Medicare FFS Beneficiaries					
	Atlantic County			10 County Rate		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change
Assessments						
• Depression Screening**	1.12	11.61	936.61	4.81	12.03	150.10
• Psychiatric Diagnostic Procedures	59.08	52.76	-10.70	53.41	45.69	-14.45
• Neuropsychological Tests	7.14	8.96	25.49	9.48	10.85	14.45
Therapy						
• Individual Psychotherapy	52.03	47.85	-8.03	54.56	53.07	-2.73
• Family Psychotherapy	2.82	1.54	-45.39	3.43	2.42	-29.45
• Group Psychotherapy	5.25	3.46	-34.10	2.98	2.71	-9.06
Psychiatric Hospital Admissions	7.19	6.55	-8.90	8.50	7.13	-16.06

* The top five risk factors were identified based on findings from a literature review (Appendix B) and factors available through Medicare claims. Logistic regression analysis was conducted with Medicare claims.

** Depression Screening comparison time frames are different (January 1, 2012 – December 31, 2012 vs. January 1, 2013 – December 31, 2013).



TOTAL MEDICARE FFS BENEFICIARY POPULATION BY COUNTY

Figure 2. Total Medicare FFS Beneficiaries by County*			
County	10/1/11-9/30/12	1/1/13-12/31/13	Absolute Change
Atlantic	47,571	46,666	-905
Bergen	142,502	139,126	-3,376
Cape May	23,769	23,291	-478
Essex	96,277	90,946	-5,331
Hudson	67,359	63,548	-3,811
Middlesex	107,061	104,272	-2,789
Monmouth	101,644	100,021	-1,623
Ocean	126,653	121,962	-4,691
Somerset	43,115	42,860	-255
Union	73,144	70,331	-2,813
10 counties**	822,505	803,020	-19,485


* Total beneficiaries who were under Medicare FFS coverage for at least one month during the time frame.

**Computing the total of all 10 counties in this table will not equal the total shown, as some beneficiaries moved from one county to another during this time frame.

The total Medicare FFS beneficiary population of Atlantic County prior to Superstorm Sandy was 47,571. After the storm, the population decreased to 46,666.

PERCENT OF MEDICARE FFS BENEFICIARIES IN THE GENERAL POPULATION

FIGURE 3. PERCENT OF MEDICARE FFS BENEFICIARIES IN THE GENERAL POPULATION IN 2012*



* Source: Medicare denominator file CY 2012, U.S. Census Bureau, American Cancer Survey (ACS), 2012 <http://www.census.gov/>.

Medicare FFS beneficiaries made up 16.72% of Atlantic County's general population in calendar year 2012.

PERCENT OF MEDICARE FFS BENEFICIARY POPULATION BY GENDER BY COUNTY

Figure 4. Percent of Medicare FFS Beneficiary Population by Female by County

County	10/1/11-9/30/12	1/1/13-12/31/13	Absolute Change*
Atlantic	55.11	54.99	-0.12
Bergen	57.00	56.73	-0.28
Cape May	53.96	53.88	-0.08
Essex	57.32	57.08	-0.23
Hudson	57.49	57.18	-0.31
Middlesex	56.31	56.06	-0.24
Monmouth	56.24	56.01	-0.23
Ocean	57.16	56.96	-0.20
Somerset	56.63	56.23	-0.40
Union	57.23	57.04	-0.19
10 counties	56.72	56.48	-0.24

* Due to rounding, the absolute change may not be the same as the difference subtracted from the two time frames shown.

Prior to Superstorm Sandy, females made up 55.11% of the entire Medicare FFS population in Atlantic County and males 44.89%. After the storm, the female beneficiary population decreased to 54.99% and males increased to 45.01%.

PERCENT OF MEDICARE FFS BENEFICIARY POPULATION BY RACE BY COUNTY

FIGURE 5. PERCENT OF MEDICARE FFS BENEFICIARY POPULATION BY RACE BY COUNTY

		County										
		Atlantic	Bergen	Cape May	Essex	Hudson	Middlesex	Monmouth	Ocean	Somerset	Union	10 Counties
White	10/1/11-9/30/12	78.73	83.77	94.56	54.95	61.88	77.80	87.88	95.60	82.96	70.08	78.96
	1/1/13-12/31/13	78.81	83.00	94.53	55.19	61.02	77.00	87.54	95.30	81.93	69.27	78.60
	Absolute Change*	0.07	-0.76	-0.03	0.23	-0.86	-0.80	-0.35	-0.30	-1.03	-0.81	-0.37
Black	10/1/11-9/30/12	14.01	4.82	3.67	35.36	12.02	7.55	7.00	1.96	6.68	19.40	11.00
	1/1/13-12/31/13	13.43	4.79	3.46	34.58	12.04	7.57	6.70	1.92	6.72	19.54	10.76
	Absolute Change*	-0.58	-0.03	-0.21	-0.78	0.02	0.02	-0.30	-0.04	0.04	0.14	-0.24
Hispanic	10/1/11-9/30/12	2.52	2.48	0.40	4.21	14.98	3.53	0.81	0.57	1.38	5.46	3.48
	1/1/13-12/31/13	2.55	2.40	0.36	4.01	14.75	3.38	0.79	0.56	1.36	5.43	3.37
	Absolute Change*	0.03	-0.07	-0.04	-0.21	-0.23	-0.15	-0.02	-0.01	-0.03	-0.03	-0.12
Asian	10/1/11-9/30/12	3.00	4.65	0.40	1.87	5.38	6.38	1.57	0.57	4.46	1.88	3.14
	1/1/13-12/31/13	3.09	4.78	0.35	1.95	5.59	6.51	1.57	0.58	4.60	1.92	3.22
	Absolute Change*	0.09	0.13	-0.05	0.08	0.21	0.12	0.00	0.01	0.13	0.03	0.08
Other	10/1/11-9/30/12	1.73	4.29	0.98	3.59	5.74	4.74	2.73	1.30	4.51	3.18	3.42
	1/1/13-12/31/13	2.12	5.03	1.31	4.27	6.61	5.54	3.40	1.65	5.40	3.85	4.06
	Absolute Change*	0.39	0.74	0.32	0.68	0.87	0.81	0.67	0.34	0.89	0.67	0.64

* Due to rounding, the absolute change may not be the same as the difference subtracted from the two time frames shown.

Both before and after Superstorm Sandy, the majority of Medicare FFS beneficiaries in Atlantic County were White followed by Black, Asian, and Hispanic.

PERCENT OF MEDICARE FFS BENEFICIARY POPULATION BY AGE BY COUNTY

FIGURE 6. PERCENT OF MEDICARE FFS BENEFICIARY POPULATION BY AGE* BY COUNTY

		County										
		Atlantic	Bergen	Cape May	Essex	Hudson	Middlesex	Monmouth	Ocean	Somerset	Union	10 Counties
<65	10/1/11-9/30/12	17.42	8.79	13.93	16.70	16.55	13.60	12.52	11.44	9.95	13.53	12.86
	1/1/13-12/31/13	17.00	8.48	13.19	16.07	16.26	13.20	12.08	11.28	9.49	13.30	12.55
	Absolute Change**	-0.42	-0.32	-0.74	-0.63	-0.29	-0.39	-0.43	-0.17	-0.45	-0.23	-0.31
65 – 74	10/1/11-9/30/12	45.11	44.84	45.69	43.97	43.81	44.33	46.43	43.25	48.32	43.60	44.73
	1/1/13-12/31/13	46.21	45.97	47.35	44.94	44.76	45.71	47.88	44.44	49.37	45.01	45.88
	Absolute Change**	1.10	1.12	1.67	0.97	0.95	1.38	1.45	1.19	1.05	1.41	1.15
75 – 84	10/1/11-9/30/12	24.90	29.20	26.86	24.89	26.27	26.78	25.92	29.02	26.56	26.32	27.05
	1/1/13-12/31/13	24.40	28.57	26.32	24.59	25.86	26.02	25.34	28.50	26.25	25.67	26.48
	Absolute Change**	-0.50	-0.63	-0.54	-0.30	-0.41	-0.75	-0.58	-0.52	-0.30	-0.65	-0.58
85 and Above	10/1/11-9/30/12	12.57	17.16	13.52	14.44	13.37	15.29	15.13	16.28	15.18	16.55	15.35
	1/1/13-12/31/13	12.39	16.99	13.13	14.40	13.12	15.06	14.70	15.78	14.89	16.02	15.09
	Absolute Change**	-0.18	-0.18	-0.38	-0.05	-0.26	-0.23	-0.43	-0.50	-0.29	-0.52	-0.26
Average Age	10/1/11-9/30/12	70.92	73.97	72.05	71.39	71.49	72.42	72.44	73.34	73.00	72.68	72.60
	1/1/13-12/31/13	70.95	73.92	72.10	71.47	71.41	72.39	72.40	73.24	72.99	72.56	72.56
	Absolute Change**	0.03	-0.05	0.06	0.08	-0.08	-0.03	-0.04	-0.11	-0.01	-0.12	-0.05

* Age calculated as end date of time frame or date of death minus birth date.


** Due to rounding, the absolute change may not be the same as the difference subtracted from the two time frames shown.

Both before and after Superstorm Sandy, the largest age group of the Medicare FFS beneficiary population in Atlantic County was between ages 65 and 74 years old followed by beneficiaries between ages 75 and 84 years old.

The average age of Medicare FFS beneficiaries in this county increased from 70.92 before the storm to 70.95 after the storm. This is also the lowest average age of Medicare FFS beneficiaries among all 10 counties.

INCOME STATUS BY COUNTY

FIGURE 7. 2012 MEDIAN HOUSEHOLD INCOME (65 YEARS AND ABOVE)



Source: U.S. Census Bureau, American Community Survey (ACS), 2012 <http://www.census.gov/>.

According to U.S. Census data from 2012, residents aged 65 and over in Atlantic County had a median household income of \$39,246. This was lower than the average income among all 10 counties.

DEMOGRAPHICS

This page left intentionally blank

PREVALENCE AND INCIDENCE

Using Medicare FFS claims data, eight behavioral health conditions were analyzed: depression or proxy disorders, depression, adjustment disorder, anxiety disorder, post-traumatic stress disorder (PTSD), alcohol or substance abuse, substance abuse alone, and suicide and intentional self-inflicted injury. These conditions were chosen based on literature review and feedback from subject matter experts.

Claims data can underestimate the real prevalence and incidence of depression in the population and individuals with depression could be diagnosed as having anxiety or adjustment disorders, as noted by the subject matter experts consulted for this project. Therefore, HQSI created a combination measure for depression (depression or proxy disorders) which includes beneficiaries who were reported for either depression, anxiety, or adjustment disorders.

The behavioral health data from January 1, 2011 to December 31, 2013 for these different measures were calculated to quantify condition occurrence:

1. The annual prevalence bar chart compares rates in two annual time frames among all 10 counties
2. New incidence in a quarter for the specified condition that was not present in the prior 12 months (Q1 2012 – Q4 2013)
3. The yearly prevalence of the condition with quarterly rolling trends to account for seasonal variation

Refer to Appendix A for measurement calculation and Appendix E for quarterly time frames and formulae.

Summary

Figure 8. Annual Prevalence of Selected Behavioral Health Conditions per 1,000 Medicare FFS Beneficiaries

County	Depression or Proxy Disorders	Depression	Anxiety Disorders	Adjustment Disorders	Alcohol or Substance Abuse	Substance Abuse Alone	PTSD	Suicide and Intentional Self-Inflicted Injury
October 1, 2011 – September 30, 2012								
Atlantic	204.91	126.73	122.86	30.28	42.98	25.26	5.32	6.47
Bergen	184.06	123.64	96.76	26.53	19.75	9.88	2.56	3.46
Cape May	208.71	125.89	123.98	24.76	40.37	20.11	5.57	4.09
Essex	184.23	119.63	88.19	36.81	38.62	22.96	3.66	4.75
Hudson	211.72	138.29	117.35	32.63	31.87	16.46	3.28	4.33
Middlesex	180.87	117.77	96.04	25.53	25.52	14.52	4.65	3.49
Monmouth	206.98	133.28	114.40	39.16	34.42	17.34	5.15	5.28
Ocean	208.85	131.66	125.55	28.88	35.33	20.67	5.86	5.33
Somerset	177.21	114.71	95.39	28.36	26.92	14.88	4.51	3.87
Union	171.83	111.46	91.54	21.85	24.24	12.13	2.48	3.44
10 counties	192.99	124.72	105.70	29.82	30.51	16.71	4.18	4.40
January 1, 2013 – December 31, 2013								
Atlantic	210.08	125.42	128.82	30.29	39.76	22.18	5.51	6.06
Bergen	192.61	127.45	106.78	27.36	23.40	11.80	2.51	3.56
Cape May	204.77	118.74	125.15	23.79	37.25	17.20	6.59	4.44
Essex	182.99	115.44	94.01	34.18	38.46	23.96	4.47	4.64
Hudson	211.02	136.51	120.44	31.64	36.71	17.31	3.48	4.64
Middlesex	184.79	118.82	103.52	25.90	28.73	15.36	5.43	3.83
Monmouth	209.21	133.07	119.46	37.29	39.88	18.07	5.84	5.17
Ocean	220.98	135.96	142.68	26.27	42.09	22.36	6.70	5.34
Somerset	181.43	116.79	102.34	29.29	31.13	16.35	5.13	4.18
Union	175.55	111.48	98.16	22.66	26.02	12.68	2.67	2.61
10 counties	197.65	125.36	113.91	29.09	33.73	17.54	4.69	4.39

Highest
 Lowest


Prevalence of the selected behavioral health conditions before and after Superstorm Sandy in the 10 counties is color coded with highest (red) and lowest (light blue) for each condition.

In the 12 months prior to Superstorm Sandy, Atlantic County had the highest prevalence rate of alcohol or substance abuse, substance abuse alone, and suicide and intentional self-inflicted injury. After the storm, Atlantic County had the highest rate of suicide and intentional self-inflicted injury despite its decrease in prevalence.

Figure 9. Percent Change of Prevalence of Selected Behavioral Health Conditions per 1,000 Medicare FFS Beneficiaries						
	Atlantic County			10 County Rate		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change
Depression or Proxy Disorders	204.91	210.08	2.52	192.99	197.65	2.41
• Depression	126.73	125.42	-1.03	124.72	125.36	0.51
• Anxiety	122.86	128.82	4.85	105.70	113.91	7.77
• Adjustment	30.28	30.29	0.03	29.82	29.09	-2.45
Alcohol or Substance Abuse	42.98	39.76	-7.49	30.51	33.73	10.55
• Substance abuse alone	25.26	22.18	-12.19	16.71	17.54	4.97
PTSD	5.32	5.51	3.57	4.18	4.69	12.20
Suicide and intentional self-inflicted injuries	6.47	6.06	-6.34	4.40	4.39	-0.23


Atlantic County experienced an increase in depression or proxy disorders, as well as in anxiety disorders, adjustment disorders, and PTSD.

FIGURE 10. QUARTERLY NEW INCIDENCE TREND OF SELECTED BEHAVIORAL HEALTH CONDITIONS: DEPRESSION OR PROXY DISORDERS* PER 1,000 MEDICARE FFS BENEFICIARIES



* Quarterly new incidence of conditions that were not diagnosed in the prior year.

FIGURE 11. QUARTERLY NEW INCIDENCE TREND* OF OTHER SELECTED BEHAVIORAL HEALTH CONDITIONS* PER 1,000 MEDICARE FFS BENEFICIARIES



* Quarterly new incidence of conditions that were not diagnosed in the prior year.

The charts above reflect quarterly trending in new incidence of the selected behavioral health conditions among Medicare FFS beneficiaries in Atlantic County.

— BEHAVIORAL HEALTH CONDITIONS —

FIGURE 12. ANNUAL PREVALENCE TREND OF SELECTED BEHAVIORAL HEALTH CONDITIONS: DEPRESSION OR PROXY DISORDERS PER 1,000 MEDICARE FFS BENEFICIARIES





FIGURE 13. ANNUAL PREVALENCE TREND OF OTHER SELECTED BEHAVIORAL HEALTH CONDITIONS PER 1,000 MEDICARE FFS BENEFICIARIES



The charts above reflect annual trending in the prevalence of the selected behavioral health conditions among Medicare FFS beneficiaries in Atlantic County.

Depression or Proxy Disorders

Figure 14. Demographics of Depression or Proxy Disorders among Medicare FFS Beneficiaries				
	10/1/11 – 9/30/12		1/1/13 – 12/31/13	
	Number of Beneficiaries	Percent (%)	Number of Beneficiaries	Percent (%)
Race				
• White	7,294	83.09	7,631	83.68
• Black	1,028	11.71	1,017	11.15
• Hispanic	249	2.84	252	2.76
• Asian	117	1.33	112	1.23
• Other	90	1.03	107	1.17
Gender				
• Males	2,886	32.88	3,097	33.96
• Females	5,892	67.12	6,022	66.04
Age				
• Below 65	2,496	28.43	2,607	28.59
• 65-74	2,838	32.33	3,035	33.28
• 75-84	2,067	23.55	2,093	22.95
• 85 and Above	1,377	15.69	1,384	15.18
Total	8,778	100.00	9,119	100.00


This table displays the number and percentage of Medicare FFS beneficiaries of each race, gender, and age diagnosed with depression or proxy disorders before and after Superstorm Sandy. There were 8,778 beneficiaries diagnosed with depression or proxy disorders in Atlantic County before the storm. This increased to 9,119 beneficiaries after the storm.

Figure 15. Demographics of Depression or Proxy Disorders Rate per 1,000 Medicare FFS Beneficiaries						
	10/1/11 – 9/30/12			1/1/13 – 12/31/13		
	Numerator	Denominator*	Rate per 1,000 Beneficiaries	Numerator	Denominator*	Rate per 1,000 Beneficiaries
Race						
• White	7,294	33,962	214.77	7,631	34,420	221.70
• Black	1,028	5,889	174.56	1,017	5,743	177.09
• Hispanic	249	1,038	239.88	252	1,083	232.69
• Asian	117	1,247	93.83	112	1,319	84.91
• Other	90	704	127.84	107	846	126.48
Gender						
• Males	2,886	19,125	150.90	3,097	19,484	158.95
• Females	5,892	23,715	248.45	6,022	23,927	251.68
Age						
• Below 65	2,496	7,160	348.60	2,607	7,146	364.82
• 65-74	2,838	19,191	147.88	3,035	19,887	152.61
• 75-84	2,067	11,131	185.70	2,093	11,034	189.69
• 85 and Above	1,377	5,358	257.00	1,384	5,344	258.98
Total	8,778	42,840	204.90	9,119	43,411	210.06

* Total eligible beneficiaries (denominator) computed after adjusting for total enrolled FFS days divided by the total measurement days in the time frame.

This table displays the rate of Medicare FFS beneficiaries per 1,000 diagnosed with depression or proxy disorders by race, gender, and age both before and after Superstorm Sandy by different demographic groups. The numerator is the number of beneficiaries with a claim for depression or proxy disorders; the denominator is the total number of beneficiaries in the county for each group. These numbers are further broken down on the following page.


FIGURE 16. DEPRESSION OR PROXY DISORDERS RATE BY RACE PER 1,000 MEDICARE FFS BENEFICIARIES



Hispanic Medicare FFS beneficiaries have the highest rate of depression or proxy disorders followed by White and Black beneficiaries.


In the 12 months prior to Superstorm Sandy, 239.88 per 1,000 Hispanic Medicare FFS beneficiaries were diagnosed with depression or proxy disorders. After the storm, this rate decreased to 232.69 per 1,000 beneficiaries.

FIGURE 17. DEPRESSION OR PROXY DISORDERS RATE BY GENDER PER 1,000 MEDICARE FFS BENEFICIARIES




Female beneficiaries have a higher rate of depression or proxy disorder compared to males. In the 12 months prior to Superstorm Sandy, 248.45 per 1,000 female Medicare FFS beneficiaries were diagnosed with depression or proxy disorders. After the storm, this rate increased to 251.68 per 1,000 female beneficiaries.

FIGURE 18. DEPRESSION OR PROXY DISORDERS RATE BY AGE GROUP PER 1,000 MEDICARE FFS BENEFICIARIES




Beneficiaries below the age of 65 have a higher rate of depression or proxy disorders. In the 12 months prior to Superstorm Sandy, 348.60 per 1,000 Medicare FFS beneficiaries below the age of 65 were diagnosed with depression or proxy disorders. After the storm, this rate increased to 364.82 per 1,000 beneficiaries.

FIGURE 19. DEPRESSION OR PROXY DISORDERS PER 1,000 MEDICARE FFS BENEFICIARIES



The prevalence rate of depression or proxy disorders in Atlantic County in the 12 months prior to Superstorm Sandy was 204.91 per 1,000 Medicare FFS beneficiaries. After the storm, this rate increased to 210.08 per 1,000 beneficiaries, reflecting a 2.52% relative increase.

FIGURE 20. QUARTERLY NEW INCIDENCE OF DEPRESSION OR PROXY DISORDERS* PER 1,000 MEDICARE FFS BENEFICIARIES




This chart reflects trending of quarterly new incidence of depression or proxy disorders among Medicare FFS beneficiaries in Atlantic County.

* Quarterly new incidences of conditions that were non-existent (not reported) in the last 12 months.

FIGURE 21. PREVALENCE OF DEPRESSION OR PROXY DISORDERS* PER 1,000 MEDICARE FFS BENEFICIARIES IN 10 COUNTIES

October 1, 2011 – September 30, 2012

January 1, 2013 – December 31, 2013




The color-coded map of New Jersey depicts prevalence of depression or proxy disorders from high (red) to low (blue) in the 10 FEMA-declared disaster counties before and after Superstorm Sandy.

* Mapped using ZIP codes of the 10 counties.

FIGURE 22. ATLANTIC COUNTY PREVALENCE OF DEPRESSION OR PROXY DISORDERS* PER 1,000 MEDICARE FFS BENEFICIARIES

October 1, 2011 – September 30, 2012

January 1, 2013 – December 31, 2013




The color-coded map of Atlantic County depicts regional variation of prevalence of depression or proxy disorders from high (red) to low (blue) before and after Superstorm Sandy.

* Mapped using ZIP codes; may not display all the city names located within the ZIP code.


Depression

FIGURE 23. DEPRESSION PER 1,000 MEDICARE FFS BENEFICIARIES



The prevalence rate of depression in Atlantic County in the 12 months prior to Superstorm Sandy was 126.73 per 1,000 Medicare FFS beneficiaries. After the storm, this rate decreased to 125.42 per 1,000 beneficiaries, reflecting a 1.03% relative decrease.

FIGURE 24. QUARTERLY NEW INCIDENCE OF DEPRESSION* PER 1,000 MEDICARE FFS BENEFICIARIES




This chart reflects trending of quarterly new incidence of depression among Medicare FFS beneficiaries in Atlantic County.

* Quarterly new incidences of conditions that were non-existent (not reported) in the last 12 months.


Anxiety Disorders

FIGURE 25. ANXIETY DISORDERS PER 1,000 MEDICARE FFS BENEFICIARIES



The prevalence rate of anxiety disorders in Atlantic County in the 12 months prior to Superstorm Sandy was 122.86 per 1,000 Medicare FFS beneficiaries. After the storm, the rate increased to 128.82 per 1,000 beneficiaries, reflecting a 4.85% relative increase.

FIGURE 26. QUARTERLY NEW INCIDENCE OF ANXIETY DISORDERS* PER 1,000 MEDICARE FFS BENEFICIARIES




This chart reflects trending of quarterly new incidence of anxiety disorders among Medicare FFS beneficiaries in Atlantic County.

* Quarterly new incidences of conditions that were non-existent (not reported) in the last 12 months.


Adjustment Disorders

FIGURE 27. ADJUSTMENT DISORDERS PER 1,000 MEDICARE FFS BENEFICIARIES



The prevalence rate of adjustment disorders in Atlantic County in the 12 months prior to Superstorm Sandy was 30.28 per 1,000 Medicare FFS beneficiaries. After the storm, the rate increased to 30.29 per 1,000 beneficiaries, reflecting a 0.03% relative increase.

FIGURE 28. QUARTERLY NEW INCIDENCE OF ADJUSTMENT DISORDERS* PER 1,000 MEDICARE FFS BENEFICIARIES




This chart reflects trending of quarterly new incidence of adjustment disorders among Medicare FFS beneficiaries in Atlantic County.

* Quarterly new incidences of conditions that were non-existent (not reported) in the last 12 months.


Post-Traumatic Stress Disorder (PTSD)

FIGURE 29. PTSD PER 1,000 MEDICARE FFS BENEFICIARIES



The prevalence rate of PTSD in Atlantic County in the 12 months prior to Superstorm Sandy was 5.32 per 1,000 Medicare FFS beneficiaries. After the storm, the rate increased to 5.51 per 1,000 beneficiaries, reflecting a 3.57% relative increase.

FIGURE 30. QUARTERLY NEW INCIDENCE OF PTSD* PER 1,000 MEDICARE FFS BENEFICIARIES




This chart reflects trending of quarterly new incidence of PTSD among Medicare FFS beneficiaries in Atlantic County.

* Quarterly new incidences of conditions that were non-existent (not reported) in the last 12 months.

Alcohol or Substance Abuse


FIGURE 31. ALCOHOL OR SUBSTANCE ABUSE PER 1,000 MEDICARE FFS BENEFICIARIES



The alcohol or substance abuse measure includes Medicare FFS beneficiaries who were reported for either alcohol abuse or substance abuse.

The prevalence rate of alcohol or substance abuse in Atlantic County in the 12 months prior to Superstorm Sandy was 42.98 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 39.76 per 1,000 beneficiaries, reflecting a 7.49% relative decrease.

FIGURE 32. QUARTERLY NEW INCIDENCE OF ALCOHOL OR SUBSTANCE ABUSE* PER 1,000 MEDICARE FFS BENEFICIARIES




This chart reflects trending of quarterly new incidence of alcohol or substance abuse among Medicare FFS beneficiaries in Atlantic County.

* Quarterly new incidences of conditions that were non-existent (not reported) in the last 12 months.


Substance Abuse Alone

FIGURE 33. SUBSTANCE ABUSE ALONE PER 1,000 MEDICARE FFS BENEFICIARIES



The prevalence rate of substance abuse alone in Atlantic County in the 12 months prior to Superstorm Sandy was 25.26 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 22.18 per 1,000 beneficiaries, reflecting a 12.19% relative decrease.

FIGURE 34. QUARTERLY NEW INCIDENCE OF SUBSTANCE ABUSE ALONE* PER 1,000 MEDICARE FFS BENEFICIARIES




This chart reflects trending of quarterly new incidence of substance abuse alone among Medicare FFS beneficiaries in Atlantic County.

* Quarterly new incidences of conditions that were non-existent (not reported) in the last 12 months.


Suicide and Intentional Self-Inflicted Injury

FIGURE 35. SUICIDE AND INTENTIONAL SELF-INFLECTED INJURY PER 1,000 MEDICARE FFS BENEFICIARIES



The prevalence rate of suicide and intentional self-inflicted injury in Atlantic County in the 12 months prior to Superstorm Sandy was 6.47 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 6.06 per 1,000 beneficiaries. This was the highest rate among all 10 counties despite a 6.34% relative decrease in prevalence.

FIGURE 36. QUARTERLY NEW INCIDENCE OF SUICIDE AND INTENTIONAL SELF-INFLECTED INJURY PER 1,000 MEDICARE FFS BENEFICIARIES



This chart reflects trending of quarterly new incidence of suicide and intentional self-inflicted injury among Medicare FFS beneficiaries in Atlantic County.

* Quarterly new incidences of conditions that were non-existent (not reported) in the last 12 months.

RISK FACTORS FOR DEPRESSION OR PROXY DISORDERS

To identify Medicare FFS beneficiaries at risk of developing depression or proxy disorders, HQSI conducted a literature review on the potential risk factors for depression or proxy disorders. Previous studies suggested that psychosocial factors, biological factors, deteriorating physical functioning, and medication side effects could increase the risk of depression or proxy disorders.

Based on the literature review and running regression models using factors available through Medicare claims data, the top five risk factors for depression or proxy disorders were identified as: Alzheimer's disease and related disorders or senile dementia, sleep disturbance, substance or alcohol abuse or tobacco use, hip/pelvic fractures, and amputations (see Appendix B).

These risk factors were reported prior to the diagnosis of depression or proxy disorders, thus indicating development of risk factors before diagnosis. The following figures show the prevalence rates for these five conditions before and after Superstorm Sandy.

Summary


Figure 37. Percent Change of Prevalence of the Top Five Risk Factors of Depression or Proxy Disorders per 1,000 Medicare FFS Beneficiaries

	Atlantic County			10 County Rate		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change
Any of the Top Five Risk Factors for Depression or Proxy Disorders	145.06	148.52	2.39	136.36	136.15	-0.15
• Substance or Alcohol Abuse or Tobacco Use	89.50	92.63	3.50	78.33	81.78	4.40
• Alzheimer's Disease and Related Disorders or Senile Dementia	33.94	33.45	-1.44	39.11	34.91	-10.74
• Sleep Disturbance	26.94	29.03	7.76	24.24	24.78	2.23
• Hip/Pelvic Fractures	6.47	6.08	-6.03	7.95	7.66	-3.65
• Amputations*	1.17	1.17	0.00	1.11	0.99	-10.81

* Rates lower than 5 per 1,000 beneficiaries.

Atlantic County experienced an increase in any on the top five risk factors for depression or proxy disorders, substance or alcohol abuse or tobacco use, and sleep disturbances.


FIGURE 38. ANNUAL PREVALENCE TREND FOR RISK FACTORS OF DEPRESSION OR PROXY DISORDERS PER 1,000 MEDICARE FFS BENEFICIARIES



This chart reflects annual trending in prevalence of the top five risk factors for depression or proxy disorders among Medicare FFS beneficiaries in Atlantic County.

ANY OF THE TOP FIVE RISK FACTORS FOR DEPRESSION OR PROXY DISORDERS

FIGURE 39. ANNUAL PREVALENCE OF ANY OF THE TOP FIVE RISK FACTORS FOR DEPRESSION OR PROXY DISORDERS PER 1,000 MEDICARE FFS BENEFICIARIES




The prevalence rate of Medicare FFS beneficiaries with any of the top five risk factors for depression or proxy disorders in Atlantic County in the 12 months prior to Superstorm Sandy was 145.06 per 1,000 beneficiaries. After the storm, the rate increased to 148.52 per 1,000 beneficiaries.

FIGURE 40. PREVALENCE OF ANY OF THE TOP FIVE RISK FACTORS FOR DEPRESSION OR PROXY DISORDERS* PER 1,000 MEDICARE FFS BENEFICIARIES IN 10 COUNTIES

October 1, 2011 – September 30, 2012

January 1, 2013 – December 31, 2013




The color-coded map of New Jersey depicts prevalence of any of the top five risk factors from high (red) to low (blue) in the 10 FEMA-declared disaster counties before and after Superstorm Sandy.

* Mapped using ZIP codes of the 10 counties.

FIGURE 41. ATLANTIC COUNTY PREVALENCE OF ANY OF THE TOP FIVE RISK FACTORS FOR DEPRESSION OR PROXY DISORDERS* PER 1,000 MEDICARE FFS BENEFICIARIES

October 1, 2011 – September 30, 2012

January 1, 2013 – December 31, 2013




The color-coded map of Atlantic County depicts regional variation of prevalence of any of the top five risk factors from high (red) to low (blue) before and after Superstorm Sandy.

* Mapped using ZIP codes; may not display all the city names located within the ZIP code.

Alzheimer's Disease and Related Disorders or Senile Dementia


FIGURE 42. ANNUAL PREVALENCE OF ALZHEIMER'S DISEASE AND RELATED DISORDERS OR SENILE DEMENTIA PER 1,000 MEDICARE FFS BENEFICIARIES



The prevalence rate of Medicare FFS beneficiaries with Alzheimer's disease and related disorders or senile dementia in Atlantic County in the 12 months prior to Superstorm Sandy was 33.94 per 1,000 beneficiaries. After the storm, the rate decreased to 33.45 per 1,000 beneficiaries.

Sleep Disturbance


FIGURE 43. ANNUAL PREVALENCE OF SLEEP DISTURBANCE PER 1,000 MEDICARE FFS BENEFICIARIES



The prevalence rate of Medicare FFS beneficiaries with sleep disturbance in Atlantic County in the 12 months prior to Superstorm Sandy was 26.94 per 1,000 beneficiaries. After the storm, the rate increased to 29.03 per 1,000 beneficiaries.

Substance or Alcohol Abuse or Tobacco Use


FIGURE 44. ANNUAL PREVALENCE OF SUBSTANCE OR ALCOHOL ABUSE OR TOBACCO USE PER 1,000 MEDICARE FFS BENEFICIARIES



The prevalence rate of Medicare FFS beneficiaries with substance or alcohol abuse or tobacco use in Atlantic County in the 12 months prior to Superstorm Sandy was 89.50 per 1,000 beneficiaries. After the storm, the rate increased to 92.63 per 1,000 beneficiaries.

Hip/Pelvic Fractures


FIGURE 45. ANNUAL PREVALENCE OF HIP/PELVIC FRACTURES PER 1,000 MEDICARE FFS BENEFICIARIES



The prevalence rate of Medicare FFS beneficiaries with hip/pelvic fractures in Atlantic County in the 12 months prior to Superstorm Sandy was 6.47 per 1,000 beneficiaries. After the storm, the rate decreased to 6.08 per 1,000 beneficiaries. This was the lowest rate among all 10 counties.

Amputations

FIGURE 46. ANNUAL PREVALENCE OF AMPUTATIONS PER 1,000 MEDICARE FFS BENEFICIARIES



The prevalence rate of Medicare FFS beneficiaries with amputations in Atlantic County in the 12 months prior to Superstorm Sandy was 1.17 per 1,000 beneficiaries. After the storm, the rate remained the same.

OUTPATIENT BEHAVIORAL HEALTH SERVICES

Assessments

Summary

Figure 47. Annual Utilization of Behavioral Health Assessment Services per 1,000 Medicare FFS Beneficiaries

County	Depression Screening*	Psychiatric Diagnostic Procedures	Neuropsychological Tests	Diagnostic Psychological Tests**	Health and Behavior Assessment/ Intervention**
October 1, 2011 – September 30, 2012					
Atlantic	1.12	59.08	7.14	3.29	1.03
Bergen	4.33	52.45	10.53	3.04	0.42
Cape May	0.65	48.08	6.13	1.77	0.70
Essex	0.83	58.52	8.31	5.55	0.53
Hudson	2.83	50.84	16.90	7.88	0.44
Middlesex	7.51	48.20	7.83	5.47	0.98
Monmouth	4.72	61.59	10.46	6.79	0.62
Ocean	9.50	54.39	9.77	3.41	0.59
Somerset	7.11	47.59	7.20	1.90	1.02
Union	3.02	46.97	6.75	2.20	0.61
10 counties	4.81	53.41	9.48	4.39	0.65

January 1, 2013 – December 31, 2013					
Atlantic	11.61	52.76	8.96	3.73	1.41
Bergen	12.04	45.55	11.00	3.20	0.35
Cape May	0.92	41.41	5.95	1.97	1.19
Essex	5.91	53.37	9.05	2.79	0.82
Hudson	9.95	45.13	17.34	8.71	0.19
Middlesex	11.39	40.15	8.94	5.38	1.25
Monmouth	13.97	54.19	10.99	6.20	0.49
Ocean	16.27	43.28	12.54	5.34	0.24
Somerset	23.76	40.21	11.45	5.03	0.87
Union	9.20	37.01	9.28	2.71	0.78
10 counties	12.03	45.69	10.85	4.61	0.66

* Depression screening comparison time frames are different (January 1, 2012 – December 31, 2012 vs. January 1, 2013 – December 31, 2013).

** Rates lower than 5 per 1,000 beneficiaries.



HQSI analyzed five behavioral health assessment services and five behavioral health therapies. Utilization of outpatient health services is color coded with lowest (red) and highest (light blue).

UTILIZATION


Figure 48. Percent Change of Behavioral Health Service Utilization – Assessments per 1,000 Medicare FFS Beneficiaries						
	Atlantic County			10 County Rate		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change
Annual Depression screening*	1.12	11.61	936.61	4.81	12.03	150.10
Psychiatric Diagnostic Procedures	59.08	52.76	-10.70	53.41	45.69	-14.45
Neuropsychological Tests	7.14	8.96	25.49	9.48	10.85	14.45
Diagnostic Psychological Tests**	3.29	3.73	13.37	4.39	4.61	5.01
Health and Behavior Assessment/ Intervention**	1.03	1.41	36.89	0.65	0.66	1.54

* Depression screening comparison time frames are different (January 1, 2012 – December 31, 2012 vs. January 1, 2013 – December 31, 2013).

** Rates lower than 5 per 1,000 beneficiaries.

Utilization of the annual depression screening benefit in Atlantic County increased from 1.12 per 1,000 Medicare FFS beneficiaries before the storm to 11.61 per 1,000 after the storm.

FIGURE 49. ANNUAL UTILIZATION TREND OF BEHAVIORAL HEALTH ASSESSMENT SERVICES PER 1,000 MEDICARE FFS BENEFICIARIES




This chart reflects annual trending in the utilization of behavioral health assessment services among Medicare FFS beneficiaries in Atlantic County.

Depression Screening

One of the long-term goals of this project is to increase the awareness and use of Medicare-covered depression screening among at-risk Medicare FFS beneficiaries residing in the 10 counties during Superstorm Sandy.


Beginning October 2011, depression screening became a Medicare-covered service. According to the CMS Screening for Depression Booklet,⁷ Medicare Part B covers an annual screening for depression of 15 minutes in length for beneficiaries in primary care settings when staff-assisted depression care supports are in place to assure accurate diagnosis, effective treatment, and follow-up. The first quarter of data in this profile for depression screening starts on January 2012 since there were only 14 claims filed for depression screening in the last quarter of 2011.

FIGURE 50. DEPRESSION SCREENING PER 1,000 MEDICARE FFS BENEFICIARIES




The rate of depression screening in Atlantic County for calendar year 2012 was 1.12 per 1,000 Medicare FFS beneficiaries. After the storm, this rate increased to 11.61 per 1,000 beneficiaries, reflecting a 936.61% relative increase, the largest increase among all 10 counties.

FIGURE 51. QUARTERLY DEPRESSION SCREENING PER 1,000 MEDICARE FFS BENEFICIARIES



This chart reflects trending of quarterly utilization of depression screening among Medicare FFS beneficiaries in Atlantic County.

FIGURE 52. DEPRESSION SCREENING* CLAIMS FOR MEDICARE FFS BENEFICIARIES




* Depression screening is a one-time benefit in 12 months.

In calendar year 2012, 97.92% of depression screening claims were filed by physicians and 2.08% were filed by nurses. After the storm, 99.35% were filed by physicians and 0.65% were filed by others.

FIGURE 53. DEPRESSION SCREENING* PER 1,000 MEDICARE FFS BENEFICIARIES IN 10 COUNTIES

January 1, 2012 – December 31, 2012

January 1, 2013 – December 31, 2013



UTILIZATION


The color-coded map of New Jersey depicts the use of depression screening from low (red) to high (blue) in the 10 FEMA-declared disaster counties before and after Superstorm Sandy.

* Mapped using ZIP codes of the 10 counties.

FIGURE 54. ATLANTIC COUNTY DEPRESSION SCREENING* PER 1,000 MEDICARE FFS BENEFICIARIES

January 1, 2012 – December 31, 2012

January 1, 2013 – December 31, 2013




The color-coded map of Atlantic County depicts regional variation in the rates of the use of the depression screening benefit from low (red) to high (blue) before and after Superstorm Sandy.

* Mapped using ZIP codes; may not display all the city names located within the ZIP code.

Diagnostic Psychological Tests

According to the CMS Mental Health Services Billing Guide, psychological testing includes psychodiagnostic assessment of emotionality, intellectual abilities, personality, and psychopathology (e.g., Minnesota Multiphasic Personality Inventory, Rorschach, or Wechsler Adult Intelligence Scale).⁸

FIGURE 55. DIAGNOSTIC PSYCHOLOGICAL TESTS PER 1,000 MEDICARE FFS BENEFICIARIES




The rate of diagnostic psychological tests in Atlantic County in the 12 months prior to Superstorm Sandy was 3.29 per 1,000 Medicare FFS beneficiaries. After the storm, this rate increased to 3.73 per 1,000 beneficiaries.

Due to these low numbers, no percent change data has been provided for this assessment.

Health and Behavior Assessment/Intervention

According to the CMS Mental Health Services Billing Guide, health and behavior assessments are used to identify the psychological, behavioral, emotional, cognitive, and social factors important to the prevention, treatment, or management of physical health problems.⁸

FIGURE 56. HEALTH AND BEHAVIOR ASSESSMENT/INTERVENTION PER 1,000 MEDICARE FFS BENEFICIARIES




The rate of health and behavior assessment/intervention in Atlantic County in the 12 months prior to Superstorm Sandy was 1.03 per 1,000 Medicare FFS beneficiaries. After the storm, this rate increased to 1.41 per 1,000 beneficiaries. This was the highest rate among all 10 counties.

Due to these low numbers, no percent change data has been provided for this assessment.

Neuropsychological Tests


According to the CMS Mental Health Services Billing Guide, neuropsychological tests are evaluations designed to determine the functional consequences of known or suspected brain injury through testing of the neurocognitive domains responsible for language, perception, memory, learning, problem solving, and adaptation.⁸

FIGURE 57. NEUROPSYCHOLOGICAL TESTS PER 1,000 MEDICARE FFS BENEFICIARIES



The rate of neuropsychological tests in Atlantic County in the 12 months prior to Superstorm Sandy was 7.14 per 1,000 Medicare FFS beneficiaries. After the storm, this rate increased to 8.96 per 1,000 beneficiaries, reflecting a 25.49% relative increase.

FIGURE 58. NEUROPSYCHOLOGICAL TESTS CLAIMS* FOR MEDICARE FFS BENEFICIARIES



* Number of claims, instead of unique beneficiaries were used in this analysis because a beneficiary can have multiple encounters for the procedure.


In the 12 months prior to Superstorm Sandy, 68.33% of neuropsychological tests claims were filed by physicians, 29.03% were filed by psychologists, 2.05% were filed by others, and 0.59% were filed by nurses.

After the storm, 84.90% of neuropsychological tests claims were filed by physicians, 13.67% were filed by psychologists, 0.82% were filed by others, and 0.61% were filed by nurses.

Psychiatric Diagnostic Procedures


According to the CMS Mental Health Services Billing Guide, psychiatric diagnostic evaluation is an integrated biopsychosocial assessment, including history, mental status, and recommendations. The evaluation may include communication with family or other sources and review of diagnostic studies.⁸

FIGURE 59. PSYCHIATRIC DIAGNOSTIC PROCEDURES PER 1,000 MEDICARE FFS BENEFICIARIES



The rate of psychiatric diagnostic procedures in Atlantic County in the 12 months prior to Superstorm Sandy was 59.08 per 1,000 Medicare FFS beneficiaries. After the storm, this rate decreased to 52.76 per 1,000 beneficiaries, reflecting a 10.70% relative decrease.

FIGURE 60. PSYCHIATRIC DIAGNOSTIC PROCEDURES CLAIMS* FOR MEDICARE FFS BENEFICIARIES



* Number of claims, instead of unique beneficiaries were used in this analysis because a beneficiary can have multiple encounters for the procedure.

In the 12 months prior to Superstorm Sandy, 50.63% of psychiatric diagnostic procedures claims were filed by physicians, 30.39% were filed by psychologists, 9.64% were filed by nurses, 8.43% were filed by social workers, and 0.91% were filed by others.

After the storm, 46.24% of psychiatric diagnostic procedures claims were filed by physicians, 34.61% were filed by psychologists, 14.16% were filed by social workers, 3.58% were filed by nurses, and 1.41% were filed by others.

Therapies

Summary

Figure 61. Annual Utilization of Behavioral Health Therapy Services per 1,000 Medicare FFS Beneficiaries					
County	Individual Psychotherapy	Family Psychotherapy*	Group Psychotherapy*	Biofeedback Therapy*	Electroconvulsive Therapy*
October 1, 2011 – September 30, 2012					
Atlantic	52.03	2.82	5.25	0.58	0.28
Bergen	62.92	4.23	2.26	0.87	0.40
Cape May	38.97	0.98	1.39	0.56	0.56
Essex	62.28	4.12	2.36	0.33	0.40
Hudson	57.04	5.06	2.42	0.22	0.27
Middlesex	53.52	3.07	5.38	1.68	0.47
Monmouth	53.59	3.56	3.23	0.26	0.61
Ocean	48.83	2.70	1.75	0.48	0.59
Somerset	57.46	3.84	4.18	0.54	0.85
Union	43.83	2.01	2.25	0.87	0.37
10 counties	54.56	3.43	2.98	0.68	0.47
January 1, 2013 – December 31, 2013					
Atlantic	47.85	1.54	3.46	0.90	0.35
Bergen	61.52	2.96	2.60	0.39	0.40
Cape May	39.21	0.92	0.64	0.41	0.73
Essex	56.97	2.43	2.81	0.44	0.34
Hudson	57.83	2.68	2.19	0.12	0.24
Middlesex	50.86	2.45	4.79	1.90	0.55
Monmouth	52.00	2.68	2.55	0.32	0.56
Ocean	52.21	2.20	1.43	0.56	0.51
Somerset	55.09	3.76	4.03	0.87	0.72
Union	40.26	1.38	2.01	0.75	0.37
10 counties	53.07	2.42	2.71	0.68	0.46

* Rates lower than 5 per 1,000 beneficiaries.

■ Lowest	■ Highest
---	---


In the 12 months prior to Superstorm Sandy, Atlantic County's most utilized behavioral health therapy service was individual psychotherapy. After the storm, despite an 8.03% relative decrease, individual psychotherapy was still the most utilized service. There was also an increase in the utilization of biofeedback and electroconvulsive therapy.

Figure 62. Percent Change of Behavioral Health Service Utilization – Therapies per 1,000 Medicare FFS Beneficiaries						
	Atlantic County			10 County Rate		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change
Individual Psychotherapy	52.03	47.85	-8.03	54.56	53.07	-2.73
Family Psychotherapy*	2.82	1.54	-45.39	3.43	2.42	-29.45
Group Psychotherapy*	5.25	3.46	-34.10	2.98	2.71	-9.06
Biofeedback Therapy*	0.58	0.90	55.17	0.68	0.68	0.00
Electroconvulsive Therapy*	0.28	0.35	25.00	0.47	0.46	-2.13

* Rates lower than 5 per 1,000 beneficiaries.

Similar to the 10 counties, Atlantic County experienced a decrease in individual, family, and group psychotherapy after Superstorm Sandy.

FIGURE 63. ANNUAL UTILIZATION TREND OF BEHAVIORAL HEALTH THERAPY SERVICES PER 1,000 MEDICARE FFS BENEFICIARIES




This chart presents annual trending in the yearly utilization of behavioral health therapies among Medicare FFS beneficiaries in Atlantic County.

Individual Psychotherapy


According to the CMS Mental Health Services Billing Guide, individual psychotherapy is the treatment of mental illness and behavioral disturbances where the physician or other qualified health professional attempts to alleviate the emotional disturbances, reverse or change maladaptive patterns of behavior, and encourage personality growth and development. This is done through the use of definitive therapeutic communication.⁸

FIGURE 64. INDIVIDUAL PSYCHOTHERAPY PER 1,000 MEDICARE FFS BENEFICIARIES



The rate of individual psychotherapy in Atlantic County in the 12 months prior to Superstorm Sandy was 52.03 per 1,000 Medicare FFS beneficiaries. After the storm, this rate decreased to 47.85 per 1,000 beneficiaries, reflecting an 8.03% relative decrease.

**FIGURE 65. INDIVIDUAL PSYCHOTHERAPY CLAIMS*
FOR MEDICARE FFS BENEFICIARIES**



* Number of claims, instead of unique beneficiaries were used in this analysis because a beneficiary can have multiple encounters for the procedure.


In the 12 months prior to Superstorm Sandy, 46.87% of individual psychotherapy claims were filed by psychologists, 30.18% were filed by social workers, 20.37% were filed by physicians, 1.46% were filed by others, and 1.12% were filed by nurses.

After the storm, 46.74% of individual psychotherapy claims were filed by psychologists, 41.84% were filed by social workers, 9.23% were filed by physicians, 1.78% were filed by others, and 0.42% were filed by nurses.

Family Psychotherapy

According to the CMS Mental Health Services Billing Guide, family psychotherapy describes the treatment of the family unit when maladaptive behaviors of family members are exacerbating the beneficiary’s mental illness or interfering with treatment. It can also be used to assist the family in addressing the maladaptive behaviors of the patient and improve treatment compliance.⁸

FIGURE 66. FAMILY PSYCHOTHERAPY PER 1,000 MEDICARE FFS BENEFICIARIES




The rate of family psychotherapy in Atlantic County in the 12 months prior to Superstorm Sandy was 2.82 per 1,000 Medicare FFS beneficiaries. After the storm, this rate decreased to 1.54 per 1,000 beneficiaries.

Due to these low numbers, no percent change data has been provided for this therapy.

Group Psychotherapy

According to the CMS Mental Health Services Billing Guide, group psychotherapy is a form of treatment where a selected group of patients are guided by a licensed psychotherapist for the purpose of helping to change maladaptive patterns which interfere with social functioning and are associated with a diagnosable psychiatric illness.⁸

FIGURE 67. GROUP PSYCHOTHERAPY PER 1,000 MEDICARE FFS BENEFICIARIES




The rate of group psychotherapy in Atlantic County in the 12 months prior to Superstorm Sandy was 5.25 per 1,000 Medicare FFS beneficiaries. After the storm, this rate decreased to 3.46 per 1,000 beneficiaries.

Due to these low numbers, no percent change data has been provided for this therapy.

Electroconvulsive Therapy

According to the CMS Mental Health Services Billing Guide, electroconvulsive therapy (ECT) is the application of electric current to the brain through scalp electrodes to induce a single seizure to produce a therapeutic effect. It is used primarily to treat major depressive disorder when antidepressant medication should not be used because it may be harmful to the patient. This type of therapy can be used for certain other clinical conditions as well.⁸

FIGURE 68. ELECTROCONVULSIVE THERAPY PER 1,000 MEDICARE FFS BENEFICIARIES




The rate of ECT in Atlantic County in the 12 months prior to Superstorm Sandy was 0.28 per 1,000 Medicare FFS beneficiaries. After the storm, this rate increased to 0.35 per 1,000 beneficiaries.

Due to these low numbers, no percent change data has been provided for this therapy.

Biofeedback Therapy

According to the CMS Mental Health Services Billing Guide, biofeedback therapy provides visual, auditory, or other evidence of the status of certain body functions so that a person can exert voluntary control over those functions, and thereby alleviate an abnormal bodily condition.⁸

FIGURE 69. BIOFEEDBACK THERAPY PER 1,000 MEDICARE FFS BENEFICIARIES



The rate of biofeedback therapy in Atlantic County in the 12 months prior to Superstorm Sandy was 0.58 per 1,000 Medicare FFS beneficiaries. After the storm, this rate increased to 0.90 per 1,000 beneficiaries.

Due to these low numbers, no percent change data has been provided for this therapy.

INPATIENT SERVICES

Summary

Inpatient services included four measures of utilization: inpatient psychiatric facilities, acute care hospital admissions, observation stays with a subsequent hospital admission, and emergency department visits with a subsequent hospital admission.

UTILIZATION

Figure 70. Annual Utilization of Inpatient Health Services per 1,000 Medicare FFS Beneficiaries

County	Psychiatric Hospital Admissions	Acute Care Hospital Admissions	Emergency Department Visits*	Observation Stays*
October 1, 2011 – September 30, 2012				
Atlantic	7.19	368.11	292.80	16.62
Bergen	9.02	287.47	217.75	6.06
Cape May	5.11	357.46	264.97	4.88
Essex	10.00	346.65	280.60	29.72
Hudson	9.84	339.25	245.61	21.52
Middlesex	7.07	296.92	237.95	15.34
Monmouth	9.44	317.02	247.54	6.43
Ocean	7.67	334.84	252.79	4.48
Somerset	8.89	278.54	216.63	4.79
Union	8.28	277.31	219.60	7.58
10 counties	8.50	315.77	244.57	11.62

January 1, 2013 – December 31, 2013				
Atlantic	6.55	330.22	263.07	20.83
Bergen	7.54	258.44	196.30	4.53
Cape May	5.10	308.15	223.86	5.97
Essex	8.13	303.36	243.59	18.54
Hudson	8.36	306.50	252.89	20.17
Middlesex	5.76	274.86	222.97	20.14
Monmouth	7.78	281.93	223.40	5.90
Ocean	6.97	293.55	220.56	15.49
Somerset	6.39	244.19	191.13	4.11
Union	6.90	256.85	200.86	6.83
10 counties	7.13	282.33	221.31	12.34

* Emergency department visits and observation stay rates were based on inpatient Part A claims only.



Utilization of inpatient health services per 1,000 Medicare FFS beneficiaries before and after Superstorm Sandy in the 10 counties is color coded with highest (red) and lowest (light blue) for each measure. These additional services were analyzed because beneficiaries with underlying behavioral health issues may seek non-behavioral health services.


In the 12 months prior to Superstorm Sandy, Atlantic County had the highest utilization of acute care hospital admissions and emergency department visits. After the storm, Atlantic County still had the highest utilization of acute care hospital visits and emergency department visits, as well as the highest utilization of observation stays.

Figure 71. Percent Change of Inpatient Health Service Utilization per 1,000 Medicare FFS Beneficiaries						
	Atlantic County			10 County Rate		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change
Psychiatric Admissions	7.19	6.55	-8.90	8.50	7.13	-16.06
Acute Care Hospital Admissions	368.11	330.22	-10.29	315.77	282.33	-10.59
Emergency Department Visits*	292.80	263.07	-10.15	244.57	221.31	-9.51
Observation Stays*	16.62	20.83	25.33	11.62	12.34	6.26

* Emergency department visits and observation stay rates were based on inpatient Part A claims only.

Atlantic County experienced an increase in observation stays and a decrease in psychiatric and acute care hospital admissions as well as in emergency department visits.


FIGURE 72. ANNUAL UTILIZATION TREND OF INPATIENT HEALTH SERVICES PER 1,000 MEDICARE FFS BENEFICIARIES



This chart reflects annual trending in the utilization of inpatient health services among Medicare FFS beneficiaries in Atlantic County.

Psychiatric Hospital Admissions

FIGURE 73. PSYCHIATRIC HOSPITAL ADMISSIONS PER 1,000 MEDICARE FFS BENEFICIARIES




In the 12 months prior to Superstorm Sandy, standalone psychiatric hospitals or distinct part psychiatric units in acute care hospitals in Atlantic County had an admissions rate of 7.19 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 6.55 per 1,000 beneficiaries, reflecting an 8.90% relative decrease.

Acute Care Hospitals

Admissions

The following data shows all-cause utilization measures and includes all Medicare FFS beneficiaries, not just beneficiaries with behavioral health conditions.

FIGURE 74. ACUTE CARE HOSPITAL ADMISSIONS PER 1,000 MEDICARE FFS BENEFICIARIES




In the 12 months prior to Superstorm Sandy, acute care hospitals in Atlantic County had an acute care admissions rate of 368.11 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 330.22 per 1,000 beneficiaries. This was the highest rate among all 10 counties despite a 10.29% relative decrease.

Observation Stays

According to the U.S. Department of Health and Human Services, observation stays are short-term treatments and assessments provided to outpatients to determine whether Medicare FFS beneficiaries require further treatment as inpatients or can be discharged.


FIGURE 75. OBSERVATION STAYS PER 1,000 MEDICARE FFS BENEFICIARIES



In the 12 months prior to Superstorm Sandy, observation stays in acute care hospitals in Atlantic County had a rate of 16.62 per 1,000 Medicare FFS beneficiaries. After the storm, the rate increased to 20.83 per 1,000 beneficiaries, the highest rate among all 10 counties. This reflects a 25.33% relative increase.

Emergency Department Visits

FIGURE 76. EMERGENCY DEPARTMENT VISITS PER 1,000 MEDICARE FFS BENEFICIARIES



In the 12 months prior to Superstorm Sandy, emergency department visits in Atlantic County had a rate of 292.80 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 263.07 per 1,000 beneficiaries. This was the highest rate among all 10 counties despite a 10.15% relative decrease.

Within 30 Days of Acute Care Hospital Discharge

Summary

Figure 77. Annual Utilization of Inpatient Health Services Within 30 Days of Discharge per 1,000 Medicare FFS Beneficiaries			
County	30-Day Hospital Readmissions	Emergency Department Visits*	Observation Stays*
October 1, 2011 – September 30, 2012			
Atlantic	75.10	96.67	8.31
Bergen	53.19	62.71	4.09
Cape May	64.06	83.85	5.95
Essex	79.02	90.77	12.02
Hudson	75.79	82.47	11.99
Middlesex	57.82	69.31	7.62
Monmouth	57.80	71.78	6.16
Ocean	62.43	78.93	7.18
Somerset	50.97	61.32	4.69
Union	50.82	61.30	5.74
10 counties	61.76	74.31	7.28
January 1, 2013 – December 31, 2013			
Atlantic	61.02	81.44	9.19
Bergen	44.29	53.65	3.88
Cape May	50.23	69.26	5.60
Essex	63.44	76.10	8.70
Hudson	65.97	78.58	10.93
Middlesex	52.02	63.65	8.37
Monmouth	46.09	61.90	5.86
Ocean	50.03	66.11	9.82
Somerset	40.27	50.98	4.23
Union	47.02	55.02	4.81
10 counties	51.37	64.36	7.16

* Emergency department visits and observation stay rates were based on both inpatient and outpatient Part A claims.

■ Highest	■ Lowest
--	---

The second set of measures is tied to utilization of services within 30 days of an acute care episode, often used as proxy indicators of care coordination, and include hospital readmissions, observation stays, and emergency department visits that occurred within 30 days of discharge. The emergency department visits is measured as with or without a subsequent hospital admission and observation stays is measured as with or without a subsequent hospital admission.


Both before and after the storm, Atlantic County had the highest utilization of emergency department visits within 30 days of discharge compared to all 10 counties. After the storm, it also experienced an increase in observation stays within 30 days of discharge.

Figure 78. Percent Change of Inpatient Health Service Utilization Within 30 Days of Discharge per 1,000 Medicare FFS Beneficiaries						
	Atlantic County			10 County Rate		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change
30-Day Hospital Readmissions	75.10	61.02	-18.75	61.76	51.37	-16.83
Emergency Department Visits*	96.67	81.44	-15.75	74.31	64.36	-13.38
Observation Stays*	8.31	9.19	10.59	7.28	7.16	-1.61

* Emergency department visits and observation stay rates were based on both inpatient and outpatient Part A claims.

Atlantic County experienced a 10.59% relative increase in observation stays within 30 days of discharge, while the 10 county rate decreased by 1.61%.


FIGURE 79. ANNUAL UTILIZATION TREND OF INPATIENT HEALTH SERVICES WITHIN 30 DAYS OF DISCHARGE PER 1,000 MEDICARE FFS BENEFICIARIES



This chart reflects annual trending in utilization of inpatient health services within 30 days of discharge among Medicare FFS beneficiaries in Atlantic County.

30-Day Hospital Readmissions


FIGURE 80. 30-DAY HOSPITAL READMISSIONS PER 1,000 MEDICARE FFS BENEFICIARIES



In the 12 months prior to Superstorm Sandy, acute care hospitals in Atlantic County had a 30-day readmission rate of 75.10 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 61.02 per 1,000 beneficiaries, reflecting an 18.75% relative decrease.

Observation Stays Within 30 Days of Discharge


FIGURE 81. OBSERVATION STAYS WITHIN 30 DAYS OF DISCHARGE PER 1,000 MEDICARE FFS BENEFICIARIES



In the 12 months prior to Superstorm Sandy, the rate of observation stays within 30 days of discharge in Atlantic County was 8.31 per 1,000 Medicare FFS beneficiaries. After the storm, the rate increased to 9.19 per 1,000 beneficiaries, reflecting a 10.59% relative increase.

Emergency Department Visits Within 30 Days of Discharge

FIGURE 82. EMERGENCY DEPARTMENT VISITS WITHIN 30 DAYS OF DISCHARGE PER 1,000 MEDICARE FFS BENEFICIARIES



In the 12 months prior to Superstorm Sandy, the rate of emergency department visits within 30 days of discharge in Atlantic County was 96.67 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 81.44 per 1,000 beneficiaries. This was the highest rate among all 10 counties despite a 15.75% relative decrease.

Other Settings

Summary

This profile also examines the utilization of home health agency, skilled nursing facility, hospice, and medical rehabilitation services. These additional services were analyzed because Medicare FFS beneficiaries with underlying behavioral health issues may seek these non-behavioral health services.

Figure 83. Annual Utilization of Other Health Services per 1,000 Medicare FFS Beneficiaries

County	Home Health Agency Services	Skilled Nursing Facility Services	Hospice Services	Medical Rehabilitation Services
October 1, 2011 – September 30, 2012				
Atlantic	98.16	64.48	28.04	14.01
Bergen	90.68	67.60	21.85	11.60
Cape May	96.11	67.91	29.92	7.80
Essex	80.94	77.28	20.20	9.81
Hudson	97.65	72.46	18.25	8.03
Middlesex	81.86	68.88	21.07	8.93
Monmouth	96.00	71.39	30.11	16.06
Ocean	100.09	76.58	30.49	22.50
Somerset	81.46	66.98	24.98	10.37
Union	84.19	69.60	20.56	9.19
10 counties	90.62	71.08	24.27	12.75
January 1, 2013 – December 31, 2013				
Atlantic	92.63	57.91	27.06	13.18
Bergen	88.45	64.69	21.19	11.06
Cape May	89.44	61.43	28.05	7.42
Essex	77.98	72.56	19.36	8.71
Hudson	88.29	72.27	15.52	6.75
Middlesex	76.38	64.56	20.92	9.25
Monmouth	88.13	65.96	27.72	15.27
Ocean	93.45	68.46	28.24	20.55
Somerset	76.40	58.33	22.84	7.62
Union	76.61	64.12	20.80	8.87
10 counties	85.02	65.99	23.01	11.86

Highest
 Lowest


Utilization of health services per 1,000 Medicare FFS beneficiaries for these settings before and after Superstorm Sandy in the 10 counties is color coded with highest (red) and lowest (light blue) for each measure.

Both before and after Superstorm Sandy, Atlantic County had the lowest utilization of skilled nursing facility services.

Figure 84. Percent Change of Other Health Services Utilization per 1,000 Medicare FFS Beneficiaries						
	Atlantic County			10 County Rate		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change
Home Health Agency	98.16	92.63	-5.63	90.62	85.02	-6.18
Skilled Nursing Facility	64.48	57.91	-10.19	71.08	65.99	-7.16
Hospice	28.04	27.06	-3.50	24.27	23.01	-5.16
Medical Rehabilitation	14.01	13.18	-5.92	12.75	11.86	-6.98

Similar to all 10 counties, Atlantic County experienced a relative decrease in the utilization of services in other health care settings.


FIGURE 85. ANNUAL UTILIZATION TREND IN OTHER HEALTH SERVICES PER 1,000 MEDICARE FFS BENEFICIARIES



This chart reflects annual trending in the utilization of other health services among Medicare FFS beneficiaries in Atlantic County.

Home Health Agency Services


FIGURE 86. HOME HEALTH AGENCY SERVICES PER 1,000 MEDICARE FFS BENEFICIARIES



In the 12 months prior to Superstorm Sandy, the utilization rate of home health agency services in Atlantic County was 98.16 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 92.63 per 1,000 beneficiaries, reflecting a 5.63% relative decrease.

Skilled Nursing Facility Services


FIGURE 87. SKILLED NURSING FACILITY SERVICES PER 1,000 MEDICARE FFS BENEFICIARIES



In the 12 months prior to Superstorm Sandy, the utilization rate of skilled nursing facility services in Atlantic County was 64.48 per 1,000 Medicare FFS beneficiaries, the lowest rate among all 10 counties. After the storm, the rate decreased to 57.91 per 1,000 beneficiaries, again the lowest rate among all 10 counties. This change also reflects a 10.19% relative decrease.

Hospice Services


FIGURE 88. HOSPICE SERVICES PER 1,000 MEDICARE FFS BENEFICIARIES



In the 12 months prior to Superstorm Sandy, the utilization rate of hospice services in Atlantic County was 28.04 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 27.06 per 1,000 beneficiaries, reflecting a 3.50% relative decrease.

Medical Rehabilitation Services

FIGURE 89. MEDICAL REHABILITATION SERVICES PER 1,000 MEDICARE FFS BENEFICIARIES



In the 12 months prior to Superstorm Sandy, the utilization rate of medical rehabilitation services in Atlantic County was 14.01 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 13.18 per 1,000 beneficiaries, reflecting a 5.92% relative decrease.

UTILIZATION

This page left intentionally blank

APPENDIX A: BEHAVIORAL HEALTH CONDITIONS

Documentation and Technical Notes

The following defines the study population, the time frames, and the exclusion and inclusion criteria:

Data Source

- New Jersey Medicare FFS Part A and Part B claims data and denominator file

Reference Time Period

- Annual prevalence for the selected behavioral health conditions comparing October 2011 – September 2012 to January 1, 2013 – December 31, 2013
- Annual prevalence trend with quarterly rolling for the selected behavioral health conditions (data starting from January 1, 2011 to December 31, 2013)
- Quarterly new incidence trend of conditions that were not existent (not reported) in the past 12 months of the selected eight behavioral health conditions (data starting from January 1, 2012 to December 31, 2013)

Mapping Tool

- QGIS Development Team, 2014, QGIS Geographic Information System. Open Source Geospatial Foundation Project. <http://qgis.osgeo.org>
- Source: ZIP code boundaries based on the 2013 U.S. Census Tiger Files

Denominator

- Denominator was the sum of all eligible Medicare FFS beneficiaries who were in the CMS denominator file during the measurement time frame
- Eligible beneficiaries were computed after adjusting for total enrolled FFS days divided by the total measurement days in the time frame
- Where Medicare FFS enrolled days > 0

Numerator

- Unique Medicare FFS beneficiaries with disease-specific inpatient or outpatient claims during the time frame
- CCW and AHRQ disease diagnosis code match (ICD-9-CM codes) Part A dgns_cd_1-25 and dgns_e_cd_1-3; Match Part B dgns_cd_1_12

Exclusions

- HMO coverage period
- Age <18 or >= 110; Age calculated as end date of time frame or date of death – birth date
- Eligible Medicare FFS days/total measurement days = 0

Resources

More information on the classification codes, requirements, and processing of the behavioral health conditions highlighted in this profile can be located at the following links:

- Buccaneer, A General Dynamics Company. Chronic Condition Data Warehouse: Additions and Access – Task Order 10 New Clinical Conditions: Requirements and Processing

APPENDIX A

- [Internet]. [unknown]: Buccaneer, A General Dynamics Company. 2013 May 22 [cited 17 Sep 2013]. Available from: https://www.ccwdata.org/cs/groups/public/documents/document/clin_cond_algo_req_proc.pdf
- Healthcare Cost and Utilization Project (H-CUP). Clinical Classifications Software (CCS) for ICD-9-CM [Internet]. Rockville (MD): Agency for Healthcare Research and Quality; Nov 2013 [15 Sep 2013]. Available from: <http://www.hcup-us.ahrq.gov/toolssoftware/ccs/ccs.jsp>

The following table shows the ICD-9-CM codes for the eight behavioral health conditions:

Behavioral Health Conditions	Numerator: Valid ICD-9-CM Codes
Depression or Proxy Disorders (Depression, Anxiety Disorders or Adjustment Disorders)	29384, 29620, 29621, 29622, 29623, 29624, 29625, 29626, 29630, 29631, 29632, 29633, 29634, 29635, 29636, 30000, 30001, 30002, 30009, 30010, 30020, 30021, 30022, 30023, 30029, 3003, 3004, 3005, 30089, 3009, 3080, 3081, 3082, 3083, 3084, 3089, 3090, 3091, 30922, 30923, 30924, 30928, 30929, 3093, 3094, 30981, 30982, 30983, 30989, 3099, 311, 3130, 3131, 31321, 31322, 3133, 31382, 31383, V790
Depression	29620, 29621, 29622, 29623, 29624, 29625, 29626, 29630, 29631, 29632, 29633, 29634, 29635, 29636, 3004, 311, V790
Anxiety Disorders	29384, 30000, 30001, 30002, 30009, 30010, 30020, 30021, 30022, 30023, 30029, 3003, , 3005, 30089, 3009, 3080, 3081, 3082, 3083, 3084, 3089, 3130, 3131, 31321, 31322, 3133, 31382, 31383
Adjustment Disorders	3090, 3091, 30922, 30923, 30924, 30928, 30929, 3093, 3094, 30981, 30982, 30983, 30989, 3099
Post-Traumatic Stress Disorder (PTSD)	30981
Alcohol or Substance Abuse	2920, 29211, 29212, 2922, 29281, 29282, 29283, 29284, 29285, 29289, 2929, 30400, 30401, 30402, 30403, 30410, 30411, 30412, 30413, 30420, 30421, 30422, 30423, 30430, 30431, 30432, 30433, 30440, 30441, 30442, 30443, 30450, 30451, 30452, 30453, 30460, 30461, 30462, 30463, 30470, 30471, 30472, 30473, 30480, 30481, 30482, 30483, 30490, 30491, 30492, 30493, 30520, 30521, 30522, 30523, 30530, 30531, 30532, 30533, 30540, 30541, 30542, 30543, 30550, 30551, 30552, 30553, 30560, 30561, 30562, 30563, 30570, 30571, 30572, 30573, 30580, 30581, 30582, 30583, 30590, 30591, 30592, 30593, 64830, 64831, 64832, 64833, 64834, 65550, 65551, 65553, 76072, 76073, 76075, 7795, 96500, 96501, 96502, 96509, V6542 Alcohol Abuse: 2910, 2911, 2912, 2913, 2914, 2915, 2918, 29181, 29182, 29189, 2919, 30300, 30301, 30302, 30303, 30390, 30391, 30392, 30393, 30500, 30501, 30502, 30503, 76071, 9800
Substance Abuse Alone	2920, 29211, 29212, 2922, 29281, 29282, 29283, 29284, 29285, 29289, 2929, 30400, 30401, 30402, 30403, 30410, 30411, 30412, 30413, 30420, 30421, 30422, 30423, 30430, 30431, 30432, 30433, 30440, 30441, 30442, 30443, 30450, 30451, 30452, 30453, 30460, 30461, 30462, 30463, 30470, 30471, 30472, 30473, 30480, 30481, 30482, 30483, 30490, 30491, 30492, 30493, 30520, 30521, 30522, 30523, 30530, 30531, 30532, 30533, 30540, 30541, 30542, 30543, 30550, 30551, 30552, 30553, 30560, 30561, 30562, 30563, 30570, 30571, 30572, 30573, 30580, 30581, 30582, 30583, 30590, 30591, 30592, 30593, 64830, 64831, 64832, 64833, 64834, 65550, 65551, 65553, 76072, 76073, 76075, 7795, 96500, 96501, 96502, 96509, V6542
Suicide and Intentional Self-Inflicted Injury	E9500, E9501, E9502, E9503, E9504, E9505, E9506, E9507, E9508, E9509, E9510, E9511, E9518, E9520, E9521, E9528, E9529, E9530, E9531, E9538, E9539, E954, E9550, E9551, E9552, E9553, E9554, E9555, E9556, E9557, E9559, E956, E9570, E9571, E9572, E9579, E9580, E9581, E9582, E9583, E9584, E9585, E9586, E9587, E9588, E9589, E959, V6284

APPENDIX B: RISK FACTORS FOR DEPRESSION OR PROXY DISORDERS

Documentation and Technical Notes

The following defines the study population, the time frame, the exclusion and inclusion criteria, and the literature review references:

Data Source

- New Jersey Medicare FFS Part A and Part B claims data and denominator file

Reference Time Period

- Annual prevalence of risk factors for depression or proxy disorders comparing October 1, 2011 – September 30, 2012 to January 1, 2013 – December 31, 2013
- Annual prevalence trend for risk factors for depression or proxy disorders consists of nine points of data with rolling quarters (starting January 1, 2011 and December 31, 2013)

Mapping Tool

- QGIS Development Team, 2014, QGIS Geographic Information System. Open Source Geospatial Foundation Project. <http://qgis.osgeo.org>
- Source: ZIP code boundaries based on the 2013 U.S. Census Tiger Files

Denominator

- Denominator was the sum of all eligible Medicare FFS beneficiaries who were in the CMS denominator file during the measurement time frame
- Eligible beneficiaries were computed after adjusting for total enrolled FFS days divided by the total measurement days in the time frame
- Where Medicare FFS enrolled days > 0

Numerator

- Unique Medicare FFS beneficiaries with disease-specific inpatient or outpatient claims during the time frame
- CCW and AHRQ disease diagnosis code match (ICD-9-CM codes) Part A dgns_cd_1-25 and dgns_e_cd_1-3; Match Part B dgns_cd_1_12

Exclusions

- HMO coverage period
- Age <18 or >= 110; Age calculated as end date of time frame or date of death – birth date
- Eligible Medicare FFS days/total measurement days = 0

Model

- Logistic Regression Models were used to determine the top five risk factors with the highest Odds Ratios (OR) ($p < 0.001$)

Resources

More information on the classification codes, requirements, and processing of the combination measure of depression or proxy disorders which includes beneficiaries reported for either depression, anxiety, or adjustment disorders can be located at the following links:

- Buccaneer, A General Dynamics Company. Chronic Condition Data Warehouse: Additions and Access – Task Order 10 New Clinical Conditions: Requirements and Processing

- [Internet]. [unknown]: Buccaneer, A General Dynamics Company. 2013 May 22 [cited 17 Sep 2013]. Available from: https://www.ccwdata.org/cs/groups/public/documents/document/clin_cond_algo_req_proc.pdf
- Healthcare Cost and Utilization Project (H-CUP). Clinical Classifications Software (CCS) for ICD-9-CM [Internet]. Rockville (MD): Agency for Healthcare Research and Quality; Nov 2013 [15 Sep 2013]. Available from: <http://www.hcup-us.ahrq.gov/toolssoftware/ccs/ccs.jsp>

Literature Review References for Risk Factors for Depression or Proxy Disorders

National Alliance on Mental Illness. Information Helpline: Depression in Older Persons Fact Sheet [Internet]. Arlington (VA): National Alliance on Mental Illness; 2009 Oct [cited 2013 Sep 17]. Available from: <http://www.nami.org/Template.cfm?Section=Helpline1&Template=/ContentManagement/ContentDisplay.cfm&ContentID=144039>

National Institute of Mental Health. Depression: Causes and Risk Factors [Internet]. Bethesda (MD): National Institute of Mental Health; 2013 Jul [cited 2013 Sep 17]. Available from: <http://nihseniorhealth.gov/depression/causesandriskfactors/01.html>

Centers for Disease Control and Prevention and National Association of Chronic Disease Directors. The State of Mental Health and Aging in America [Internet]. Atlanta (GA): National Association of Chronic Disease Directors, 2008 [cited 2013 Sep 19]. 11 p. Available from: http://www.cdc.gov/aging/pdf/mental_health.pdf

Jacques L, Jensen T, Schafer J, Caplan S, Schott L. Final Coverage Decision Memorandum for Screening for Depression in Adults [Internet]. Baltimore (MD): Centers for Medicare & Medicaid Services; 2011 Oct 14 [cited 2013 Sep 18]. 42 p. Available from: <http://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?NCAId=251>

Thakur M, Blazer DG. Depression in long-term care. *Journal of the American Medical Directors Association* [Internet]. 2008 Feb [cited 2013 Sep 19];9(2):82-87. Available from: <http://www.amda.com/tools/clinical/depression/DepressioninLongTermCare.pdf>

Sozeri-Varma G. Depression in the elderly: clinical features and risk factors. *Aging and Disease* [Internet]. 2012 Dec [cited 2013 Sep 18];3(6):465-471. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3522513/>

Qian J, Simoni-Wastila L, Rattinger GB, Lehmann S, Langenberg P, et al. Associations of depression diagnosis and antidepressant treatment with mortality among young and disabled Medicare beneficiaries with COPD. *General Hospital Psychiatry*. 2013 Jul 18 [cited 2013 Sep 22]; 35(6):612-618.

Shao W, Ahmad R, Khutoryansky N, Aagren M, Bouchard J. Evidence supporting an association between hypoglycemic events and depression. *Current Medical Research and Opinion*. 2013 Sep 23 [cited 2013 Sep 22]: 1-7.

Substance Abuse and Mental Health Services Administration. The Treatment of Depression in Older Adults: Depression and Older Adults: Key Issues [Internet]. Rockville, MD: Center for Mental Health Services, Substance Abuse and Mental Health Services Administration, U.S. Department of Health and Human Services, 2011 [cited 2013 Sep 24]. HHS Pub. No. SMA-11-4631. 24 p. Available from: <http://store.samhsa.gov/shin/content/SMA11-4631CD-DVD/SMA11-4631CD-DVD-KeyIssues.pdf>

Himelhoch S, Weller WE, Wu AW, Anderson GF, Cooper LA. Chronic medical illness, depression, and use of acute medical services among Medicare beneficiaries. *Medical Care*. 2004 Jun [cited 2013 Sep 25];42(6):512-521.

Mohile SG, Fan L, Reeve E, Jean-Pierre P, Mustian K, et al. Association of cancer with geriatric syndromes in older Medicare beneficiaries. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology* [Internet]. 2011 Apr 10 [cited 2013 Sep 25];29(11):1458-1464. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3082984/>

Jayadevappa R, Malkowicz SB, Chhatre S, Johnson JC, Gallo JJ. The burden of depression in prostate cancer. *Psycho-oncology*. 2012 Dec [cited 2013 Sep 26];21(12):1338-1345.

Missouri Department of Mental Health. CPS Facts: Depression and Older Adults [Internet]. Jefferson City(MO): Missouri Department of Mental Health, [date unknown, cited 2013 Sep 26], 2 p. Available from: <http://dmh.mo.gov/docs/mentalillness/elderlydepress.pdf>

Oregon State University, Washington State University, University of Idaho. Depression in Later Life: Recognition and Treatment [Internet]. Corvallis(OR): Pacific Northwest Extension Publication; 2004 Jul [Published April 1990; revised July 2000; cited 2013 Sep 29]; 32 p. Available from: <http://ir.library.oregonstate.edu/xmlui/bitstream/handle/1957/20713/pnw347.pdf>

Cole MG, Dendukuri N. Risk factors for depression among elderly community subjects: a systematic review and meta-analysis. *American Journal of Psychiatry* [Internet]. 2003 Jun [cited 2013 Sep 29]; 160(6):1147-1156. Available from: <http://ajp.psychiatryonline.org/article.aspx?articleid=176272>

Kohn R, Levav I, Garcia ID, Machuca ME, Tamashiro R. Prevalence, risk factors and aging vulnerability for psychopathology following a natural disaster in a developing country. *International Journal of Geriatric Psychiatry*. 2005 Sep [cited 2013 Sep 29];20(9):835-841.

Pietrzak RH, Southwick SM, Tracy M, Galea S, Norris FH. Posttraumatic stress disorder, depression, and perceived needs for psychological care in older persons affected by Hurricane Ike. *Journal of Affective Disorders* [Internet]. 2012 Apr [cited 2013 Sep 30];138(1-2):96-103. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3306486/>

Oriol W. Psychosocial Issues for Older Adults in Disasters [Internet]. Washington (DC): Emergency Services and Disaster Relief Branch, Center for Mental Health Services (CMHS), Substance Abuse and Mental Health Services Administration; 1999 [cited 2013 Sep 30]; DHHS Publication No. ESDRB SMA 99-3323. 79 p. Available from: <http://store.samhsa.gov/shin/content/SMA99-3323/SMA99-3323.pdf>

O'Connor EA, Whitlock EP, Gaynes B, Beil TL. Screening for Depression in Adults and Older Adults in Primary Care: An Updated Systematic Review. [Internet]. Rockville (MD): Agency for Healthcare Research and Quality (US); 2009 Dec [cited 2013 Sept 30]. 167 p. (Evidence Synthesis No. 75. AHRQ Publication No. 10-05143-EF-1). Available from: <http://www.ncbi.nlm.nih.gov/books/NBK36403/pdf/TOC.pdf>

Noyes K, Liu H, Lyness JM, Friedman B. Medicare beneficiaries with depression: comparing diagnoses in claims data with the results of screening. *Psychiatric Services* [Internet]. 2011 Oct [cited 2013 Sep 30];62(10):1159-1166. Available from: http://ps.psychiatryonline.org/data/Journals/PSS/4336/pss6210_1159.pdf

The following table shows the ICD-9-CM codes for the top five risk factors for depression or proxy disorders:

Top Five Risk Factors for Depression or Proxy Disorders*	Numerator: Valid ICD-9-CM Codes
Alzheimer's Disease and Related Disorders or Senile Dementia	3311, 33111, 33119, 3312, 3317, 2900, 29010, 29011, 29012, 29013, 29020, 29021, 2903, 29040, 29041, 29042, 29043, 2940, 2941, 29410, 29411, 2948, 797
Sleep Disturbance	04672, 29182, 29285, 30740, 30741, 30742, 30748, 30749, 32700, 32701, 32702, 32709, 78050, 78051, 78052, 78059
Substance or Alcohol Abuse or Tobacco Use	2910, 2911, 2912, 2913, 2914, 2915, 2918, 29181, 29182, 29189, 2919, 2920, 29211, 29212, 2922, 29281, 29282, 29283, 29284, 29285, 29289, 2929, 30300, 30301, 30302, 30303, 30390, 30391, 30392, 30393, 30400, 30401, 30402, 30403, 30410, 30411, 30412, 30413, 30420, 30421, 30422, 30423, 30430, 30431, 30432, 30433, 30440, 30441, 30442, 30443, 30450, 30451, 30452, 30453, 30460, 30461, 30462, 30463, 30470, 30471, 30472, 30473, 30480, 30481, 30482, 30483, 30490, 30491, 30492, 30493, 30500, 30501, 30502, 30503, 3051, 30510, 30511, 30512, 30513, 30520, 30521, 30522, 30523, 30530, 30531, 30532, 30533, 30540, 30541, 30542, 30543, 30550, 30551, 30552, 30553, 30560, 30561, 30562, 30563, 30570, 30571, 30572, 30573, 30580, 30581, 30582, 30583, 30590, 30591, 30592, 30593, 33392, 3575, 4255, 5353, 53530, 53531, 5710, 5711, 5712, 5713, 64830, 64831, 64832, 64833, 64834, 65550, 65551, 65553, 76071, 76072, 76073, 76075, 7795, 7903, 96500, 96501, 96502, 96509, 9800, V110, V111, V112, V113, V114, V118, V119, V154, V1541, V1542, V1549, V1582, V6285, V6542, V663, V701, V702, V7101, V7102, V7109, V790, V791, V792, V793, V798, V799
Hip/Pelvic Fractures	73314, 73315, 73396, 73397, 73398, 8080, 8081, 8082, 8083, 80841, 80842, 80843, 80849, 80851, 80852, 80853, 80859, 8088, 8089, 82000, 82001, 82002, 82003, 82009, 82010, 82011, 82012, 82013, 82019, 82020, 82021, 82022, 82030, 82031, 82032, 8208, 8209
Amputations	8870, 8871, 8872, 8873, 8874, 8875, 8876, 8877, 8960, 8961, 8962, 8963, 8970, 8971, 8972, 8973, 8974, 8975, 8976, 8977, 9059, 99760, 99761, 99762, 99769

* Other risk factors for depression or proxy disorders analyzed include Acute Myocardial Infarction (AMI), Stroke/Transient Ischemic Attack, Coronary Artery Bypass Graft Surgery (CABG), Parkinson's Disease, Chronic Obstructive Pulmonary Disease and Bronchiectasis (COPD), Diabetes, Chronic Kidney Disease, Rheumatoid Arthritis/Osteoarthritis (RA/OA), Macular Degeneration, Disability, History of Cancer, Heart Failure, and Acquired Hypothyroidism.

APPENDIX C: UTILIZATION OF OUTPATIENT MENTAL HEALTH SERVICES

Documentation and Technical Notes

The following defines the study population, the time frame, and the exclusion and inclusion criteria:

Data Source

- New Jersey Medicare FFS Part A and Part B claims data and denominator file

Reference Time Period

- Annual utilization comparing October 1, 2011 – September 30, 2012 to January 1, 2013 – December 31, 2013
- Annual utilization trend consists of nine points of data with rolling quarters (starting January 1, 2011 and ending December 31, 2013)
- Quarterly utilization trend charts for depression screening contains data from January 1, 2012 to December 31, 2013

Mapping Tool

- QGIS Development Team, 2014, QGIS Geographic Information System. Open Source Geospatial Foundation Project. <http://qgis.osgeo.org>
- Source: ZIP code boundaries based on the 2013 U.S. Census Tiger Files

Denominator

- Denominator was the sum of all eligible Medicare FFS beneficiaries who were in the CMS denominator file during the measurement time frame
- Eligible beneficiaries were computed after adjusting for total enrolled FFS days divided by the total measurement days in the time frame
- Where Medicare FFS enrolled days > 0

Numerator

Unique Medicare FFS beneficiaries with specific outpatient mental health service claims

Exclusions

- HMO coverage period
- Age <18 or >= 110; Age calculated as end date of time frame or date of death – birth date
- Eligible Medicare FFS days/total measurement days =0

Resources

More information on the definitions and uses of the outpatient mental health services highlighted in this profile can be located at http://www.cmsbilling.org/forms/NHIC_Medicare_B_Mental_Health_billing_guide_2008.pdf.

APPENDIX C

The following table shows the CPT/HCPCS codes for the outpatient mental health services:

Mental Health Services	Numerator: CPT/HCPCS Codes
Assessments	
Depression Screening	G0444
Diagnostic Psychological Tests	96101, 96102, 96103, 96105, 96110, 96111
Health and Behavior Assessment/Intervention	96150, 96151, 96152, 96153, 96154, 96155
Neuropsychological Tests	96116, 96118, 96119, 96120
Psychiatric Diagnostic Procedures	90801, 90802, 90791, 90792
Therapies	
Individual Psychotherapy	90804, 90805, 90832, 90833, 90806, 90807, 90834, 90836, 90808, 90809, 90810, 90811, 90812, 90813, 90814, 90815, 90816, 90817, 90818, 90819, 90821, 90822, 90823, 90824, 90826, 90827, 90828, 90829, 90837, 90838, 90839, 90840
Family Psychotherapy	90846, 90847
Group Psychotherapy	90849, 90853, 90857
Electroconvulsive Therapy	90870
Biofeedback Therapy	90901, 90911

APPENDIX D: UTILIZATION OF SERVICES – INPATIENT AND OTHER SETTINGS

Documentation and Technical Notes

The following defines the study population, the time frame, and the exclusion and inclusion criteria:

Data Source

New Jersey Medicare FFS Part A claims data and denominator file

Reference Time Period

- Annual utilization comparing October 1, 2011 – September 30, 2012 to January 1, 2013 – December 31, 2013
- Annual utilization trend consists of nine points of data with rolling quarters (starting January 1, 2011 and ending December 31, 2013)

Denominator

- Denominator was the sum of all eligible Medicare FFS beneficiaries who were in the CMS denominator file during the measurement time frame
- Eligible beneficiaries were computed after adjusting for total enrolled FFS days divided by the total measurement days in the time frame
- Where Medicare FFS enrolled days > 0

Exclusions

- HMO coverage period
- Age <18 or \geq 110; Age calculated as end date of time frame or date of death – birth date
- Eligible Medicare FFS days/total measurement days =0

Utilization Measure

Refer to Appendix E.

Numerator

Utilization Measure Description	Numerator
Acute Care Hospital Admission	Number of inpatient admissions (Nch_clm_type_cd = 60, 61)
30-Day Hospital Readmissions	Number of readmissions that occurred within 30 days of hospital discharge (Nch_clm_type_cd = 60, 61)
Emergency Department Visits	Number of emergency department visits, with subsequent inpatient admission (Nch_clm_type_cd = 60, 61 and revenue code in '0450''0451''0452''0456''0459''0981')
Emergency Department Visits within 30 Days of Hospital Discharge	Number of emergency department visits within 30 days of hospital discharge, with or without subsequent admission (Nch_clm_type_cd = 60, 61, 40 and revenue code in '0450''0451''0452''0456''0459''0981')
Observation Stays	Number of observation stays, with subsequent inpatient admission (Nch_clm_type_cd = 60, 61 and revenue code in '0762')
Observation Stays within 30 Days of Hospital Discharge	Number of observation stays within 30 days of hospital discharge, with or without subsequent admission (Nch_clm_type_cd = 60, 61, 40 and revenue code in '0762')
Home Health Agency Services	Number of eligible beneficiaries with at least one home health agency claim (Nch_clm_type_cd = 10)
Skilled Nursing Facility Services	Number of eligible beneficiaries with at least one skilled nursing facility claim (Nch_clm_type_cd = 20, 30)
Hospice Services	Number of eligible beneficiaries with at least one hospice claim (Nch_clm_type_cd = 50)
Medical Rehabilitation Services	Number of eligible beneficiaries with at least one medical rehabilitation claim (Nch_clm_type_cd = 60, 61 and hsp_id format: xxTxxx or between xx3025 and xx3099)
Psychiatric Hospital Admissions	Number of eligible beneficiaries with at least one psychiatric hospital admission claim (Nch_clm_type_cd = 60, 61 and hsp_id format: xxSxxx or between xx4000 and xx4499)

APPENDIX E: TIME FRAMES AND FORMULAE

Time Frames	
Quarters	Dates
Q1	January 1 to March 31
Q2	April 1 to June 30
Q3	July 1 to September 30
Q4	October 1 to December 31

Formulae

$$\text{Incidence} = \frac{\text{(Number of unique beneficiaries with new cases during the time frame, condition not present in the past 12 months)}}{\text{(Total unique beneficiaries in the population during the time frame)}}$$

$$\text{Prevalence} = \frac{\text{(Number of unique beneficiaries with the condition during the time frame)}}{\text{(Total unique beneficiaries in the population during the time frame)}}$$

$$\text{Utilization} = \frac{\text{(Number of unique beneficiaries or measures with specific service utilization)}}{\text{(Total unique beneficiaries in the population during the time frame)}}$$

$$\text{Relative change} = \frac{\text{(Current rate-Former rate)}}{\text{(Former rate)}}$$

APPENDIX F: PROFESSIONAL TYPE BY BEHAVIORAL HEALTH SERVICES

The following defines the data source and time period for the provider summary tables and listings:

Data Source

New Jersey Medicare FFS Part B claims data

Reference Time Period

- Professional type of behavioral health service claims during October 1, 2011 – September 30, 2012 and January 1, 2013 – December 31, 2013

Professional Type Credentials

- Physicians: DO, MD
- Psychologists: PhD, PsyD, EdD
- Social Workers: MSW, LCSW
- Nurses: APN, RN, NP
- Others: Other

APPENDIX G: REFERENCES

1. Weiss MG, Saraceno B, Saxena S, van Ommeren M. Mental health in the aftermath of disasters: consensus and controversy. *The Journal of Nervous and Mental Disease*. 2003 Sep; 191(9):611-615.
2. Foa EB, Stein DJ, McFarlane AC. Symptomatology and psychopathology of mental health problems after disaster. *The Journal of Clinical Psychology* [Internet]. 2006;[cited 16 Sep 2013];67 Suppl 2:15-25.
3. Wang PS, Gruber MJ, Powers RE, Schoenbaum M, Speier AH, Wells KB, Kessler RC. Mental health service use among hurricane Katrina survivors in the eight months after the disaster. *Psychiatry Services* [Internet]. 2007 Nov [cited 16 Sep 2013]; 58(11):1403-1411. Available from: <http://ps.psychiatryonline.org/data/Journals/PSS/3824/07ps1403.pdf>
4. Voelker R. Post-katrina mental health needs prompt group to compile disaster medicine guide. *JAMA*. 2006 Jan [cited 2013 Sep 17]; 295(3):259-260.
5. Centers for Medicare & Medicaid Services. Medicare Claims Database, Parts A and B, January 1, 2011 – March 31, 2013. Baltimore (MD): CMS, Department of Health and Human Services. Accessed: September 15, 2013.
6. U.S. Department of Commerce: United States Census Bureau, American Fact Finder [Internet]. Washington (DC): U.S. Department of Commerce. Median Income in the Past 12 Months (in 2012 Inflation-Adjusted Dollars); 2012 [cited 15 Sep 2013]; [about 2 screens]. Available from: http://factfinder2.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_12_1YR_S1903&prodType=table
7. Centers for Medicare & Medicaid Services. Screening for Depression, February 2013. Available from: <https://www.cms.gov/Outreach-and-Education/Medicare-Learning-Network-MLN-MLNProducts/Downloads/Screening-for-Depression-Booklet-ICN907799.pdf>
8. Centers for Medicare & Medicaid Services. Mental Health Services Billing Guide, April 2013. Hingham (MA): NHIC, Corp. Apr 2013. 40 p.

APPENDIX G

This page left intentionally blank

Executive Summary

Figure 1. Snapshot of Atlantic County	10
---------------------------------------	----

Demographics

Figure 2. Total Medicare FFS Beneficiaries by County	11
Figure 3. Percent of Medicare FFS Beneficiaries in the General Population in 2012	12
Figure 4. Percent of Medicare FFS Beneficiary Population by Female by County	12
Figure 5. Percent of Medicare FFS Beneficiary Population by Race by County	13
Figure 6. Percent of Medicare FFS Beneficiary Population by Age by County	14
Figure 7. 2012 Median Household Income (65 years and above)	15

Prevalence and Incidence

Figure 8. Annual Prevalence of Selected Behavioral Health Conditions	18
Figure 9. Percent Change of Prevalence of Selected Behavioral Health Conditions	19
Figure 10. Quarterly New Incidence Trend of Selected Behavioral Health Conditions: Depression or Proxy Disorders	19
Figure 11. Quarterly New Incidence Trend of Other Selected Behavioral Health Conditions	19
Figure 12. Annual Prevalence Trend of Selected Behavioral Health Conditions: Depression or Proxy Disorders	20
Figure 13. Annual Prevalence Trend of Other Selected Behavioral Health Conditions	20
Figure 14. Demographics of Depression or Proxy Disorders	21
Figure 15. Demographics of Depression or Proxy Disorders Rate	22
Figure 16. Depression or Proxy Disorders Rate by Race	23
Figure 17. Depression or Proxy Disorders Rate by Gender	23
Figure 18. Depression or Proxy Disorders Rate by Age Group	23
Figure 19. Depression or Proxy Disorders (Annual Prevalence, Annual Trend, and Percent Change)	24
Figure 20. Quarterly New Incidence of Depression or Proxy Disorders	24
Figure 21. Prevalence of Depression or Proxy Disorders in 10 Counties	25
Figure 22. Atlantic County Prevalence of Depression or Proxy Disorders	26
Figure 23. Depression (Annual Prevalence, Annual Trend, and Percent Change)	27
Figure 24. Quarterly New Incidence of Depression	27
Figure 25. Anxiety Disorders (Annual Prevalence, Annual Trend, and Percent Change)	28
Figure 26. Quarterly New Incidence of Anxiety Disorders	28
Figure 27. Adjustment Disorders (Annual Prevalence, Annual Trend, and Percent Change)	29
Figure 28. Quarterly New Incidence of Adjustment Disorders	29
Figure 29. PTSD (Annual Prevalence, Annual Trend, and Percent Change)	30
Figure 30. Quarterly New Incidence of PTSD	30
Figure 31. Alcohol or Substance Abuse (Annual Prevalence, Annual Trend, and Percent Change)	31
Figure 32. Quarterly New Incidence of Alcohol or Substance Abuse	31
Figure 33. Substance Abuse Alone (Annual Prevalence, Annual Trend, and Percent Change)	32
Figure 34. Quarterly New Incidence of Substance Abuse Alone	32
Figure 35. Suicide and Intentional Self-Inflicted Injury (Annual Prevalence, Annual Trend, and Percent Change)	33
Figure 36. Quarterly New Incidence of Suicide and Intentional Self-Inflicted Injury	33

Risk Factors for Depression or Proxy Disorders

Figure 37. Percent Change of Prevalence of the Top Five Risk Factors of Depression or Proxy Disorders 34

Figure 38. Annual Prevalence Trend for Risk Factors of Depression or Proxy Disorders 35

Figure 39. Annual Prevalence of Any of the Top Five Risk Factors for Depression or Proxy Disorders 35

Figure 40. Prevalence of Any of the Top Five Risk Factors for Depression or Proxy Disorders in 10 Counties 36

Figure 41. Atlantic County Prevalence of Any of the Top Five Risk Factors for Depression or Proxy Disorders 37

Figure 42. Annual Prevalence of Alzheimer's Disease and Related Disorders or Senile Dementia 38

Figure 43. Annual Prevalence of Sleep Disturbance 38

Figure 44. Annual Prevalence of Substance or Alcohol Abuse or Tobacco Use 39

Figure 45. Annual Prevalence of Hip/Pelvic Fractures 39

Figure 46. Annual Prevalence of Amputations 40

Outpatient Behavioral Health Services

Assessments

Figure 47. Annual Utilization of Behavioral Health Assessment Services 41

Figure 48. Percent Change of Behavioral Health Service Utilization – Assessments 42

Figure 49. Annual Utilization Trend of Behavioral Health Assessment Services 42

Figure 50. Depression Screening (Annual Utilization, Annual Trend, and Percent Change) 43

Figure 51. Quarterly Depression Screening 44

Figure 52. Depression Screening Claims for Medicare FFS Beneficiaries 44

Figure 53. Depression Screening in 10 Counties 45

Figure 54. Atlantic County Depression Screening 46

Figure 55. Diagnostic Psychological Tests (Annual Utilization and Annual Trend) 47

Figure 56. Health and Behavior Assessment/Intervention (Annual Utilization and Annual Trend) 48

Figure 57. Neuropsychological Tests (Annual Utilization, Annual Trend, and Percent Change) 49

Figure 58. Neuropsychological Tests Claims for Medicare FFS Beneficiaries 50

Figure 59. Psychiatric Diagnostic Procedures (Annual Utilization, Annual Trend, and Percent Change) 51

Figure 60. Psychiatric Diagnostic Procedures Claims for Medicare FFS Beneficiaries 52

Therapies

Figure 61. Annual Utilization of Behavioral Health Therapy Services 53

Figure 62. Percent Change of Behavioral Health Service Utilization – Therapies 54

Figure 63. Annual Utilization Trend of Behavioral Health Therapy Services 54

Figure 64. Individual Psychotherapy (Annual Utilization, Annual Trend, and Percent Change) 55

Figure 65. Individual Psychotherapy Claims for Medicare FFS Beneficiaries 56

Figure 66. Family Psychotherapy (Annual Utilization and Annual Trend)	57
Figure 67. Group Psychotherapy (Annual Utilization and Annual Trend)	58
Figure 68. Electroconvulsive Therapy (Annual Utilization and Annual Trend)	59
Figure 69. Biofeedback Therapy (Annual Utilization and Annual Trend)	60
Inpatient Services	
Figure 70. Annual Utilization of Inpatient Health Services	61
Figure 71. Percent Change of Inpatient Health Service Utilization	62
Figure 72. Annual Utilization Trend of Inpatient Health Services	62
Figure 73. Psychiatric Hospital Admissions (Annual Utilization, Annual Trend, and Percent Change)	63
<i>Acute Care Hospitals</i>	
Figure 74. Acute Care Hospital Admissions (Annual Utilization, Annual Trend, and Percent Change)	64
Figure 75. Observation Stays (Annual Utilization, Annual Trend, and Percent Change)	65
Figure 76. Emergency Department Visits (Annual Utilization, Annual Trend, and Percent Change)	66
<i>Within 30 Days of Acute Care Hospital Discharge</i>	
Figure 77. Annual Utilization of Inpatient Health Services Within 30 Days of Discharge	67
Figure 78. Percent Change of Inpatient Health Service Utilization Within 30 Days of Discharge	68
Figure 79. Annual Utilization Trend of Inpatient Health Services Within 30 Days of Discharge	68
Figure 80. 30-Day Hospital Readmissions (Annual Utilization, Annual Trend, and Percent Change)	69
Figure 81. Observation Stays Within 30 Days of Discharge (Annual Utilization, Annual Trend, and Percent Change)	70
Figure 82. Emergency Department Visits Within 30 Days of Discharge (Annual Utilization, Annual Trend, and Percent Change)	71
<i>Other Settings</i>	
Figure 83. Annual Utilization of Other Health Services	72
Figure 84. Percent Change of Other Health Services Utilization	73
Figure 85. Annual Utilization Trend in Other Health Services	73
Figure 86. Home Health Agency Services (Annual Utilization, Annual Trend, and Percent Change)	74
Figure 87. Skilled Nursing Facility Services (Annual Utilization, Annual Trend, and Percent Change)	75
Figure 88. Hospice Services (Annual Utilization, Annual Trend, and Percent Change)	76
Figure 89. Medical Rehabilitation Services (Annual Utilization, Annual Trend, and Percent Change)	77

INDEX OF FIGURES

This page left intentionally blank



**Quality Improvement
Organizations**

Sharing Knowledge. Improving Health Care.
CENTERS FOR MEDICARE & MEDICAID SERVICES



557 Cranbury Road, Suite 21
East Brunswick, NJ 08816
T 732-238-5570 • F 732-238-7766
www.hqsi.org